

The limits of the total crystal-field splittings

J. Mulak¹, M. Mulak²

¹ *Trzebiatowski Institute of Low Temperature and Structure Research,
Polish Academy of Sciences, 50-950, PO Box 1410, Wroclaw, Poland*

² *Institute of Physics, Wroclaw University of Technology,
Wyb. Wyspianskiego 27, 50-370 Wroclaw, Poland*

Corresponding author:

Prof. Jacek Mulak

Trzebiatowski Institute of Low Temperature and Structure Research,
Polish Academy of Sciences, 50-950, PO Box 1410, Wroclaw, POLAND

email: Maciej.Mulak@pwr.wroc.pl

Tel: (+4871) 3435021, 3443206

Fax: (+4871) 3441029

Abstract

The crystal-fields causing $|J\rangle$ electron states splittings of the same second moment σ^2 can produce different total splittings ΔE magnitudes. Based on the numerical data on crystal-field splittings for the representative sets of crystal-field Hamiltonians $\mathcal{H}_{\text{CF}} = \sum_k \sum_q B_{kq} C_q^{(k)}$ with fixed indexes either k or q , the potentials leading to the extreme ΔE have been identified. For all crystal-fields the admissible ranges $(\Delta E_{\min}, \Delta E_{\max})$ have been found numerically for $1 \leq J \leq 8$. The extreme splittings are reached in the crystal-fields for which $\mathcal{H}_{\text{CF}}$ s are the definite superpositions of the $C_q^{(k)}$ components with different rank $k = 2, 4$ and 6 and the same index q . Apart from few exceptions, the lower limits $\Delta E_{\min}$ occur in the axial fields of $\mathcal{H}_{\text{CF}}(q=0) = B_{20}C_0^{(2)} + B_{40}C_0^{(4)} + B_{60}C_0^{(6)}$, whereas the upper limits $\Delta E_{\max}$ in the low symmetry fields of $\mathcal{H}_{\text{CF}}(q=1) = B_{21}C_1^{(2)} + B_{41}C_1^{(4)} + B_{61}C_1^{(6)}$. Mixing the $\mathcal{H}_{\text{CF}}$ components with different q yields a secondary effect and does not determine the extreme splittings. The admissible $\Delta E_{\min}$ changes with J from 2.00σ to 2.40σ , whereas the $\Delta E_{\max}$ from 2.00σ to 4.10σ . The maximal gap $\Delta E_{\max} - \Delta E_{\min} = 2.00\sigma$ has been found for the states $|J=4\rangle$. Not all the nominally allowed total splittings, preserving $\sigma^2 = \text{const}$ condition, are physically available, and in consequence not all virtual splittings diagrams can be observed in real crystal-fields.

PACS: 71.15.-m, 71.23.An, 71.70.Ch, 75.10.Dg

1. Introduction

The total splitting ranges $(\Delta E_{min}, \Delta E_{max})$ of free ion electron states in different axial crystal-fields (CFs) yielding the splittings of the same second moment σ^2 have been thoroughly analyzed in our previous paper [1]. Here, we extend this analysis for all CFs. Throughout the paper the tensor Wybourne notation [2] for the CF Hamiltonian $\mathcal{H}_{CF}$ and the CF parameters B_{kq} (CFP) is consistently used (Eq.(1)),

$$\mathcal{H}_{CF} = \sum_k \sum_q B_{kq} C_q^{(k)}. \quad (1)$$

The zero-approximation initial states forming the interaction matrices are the Russell-Saunders coupled states $|\alpha SLJ\rangle$ of well defined quantum number J with the degeneracy $2J + 1$, where α stands for the remaining needed quantum numbers. For the sake of simplicity the initial states are denoted as $|J\rangle$. Further extension of the studies for the mixed states as well as the states resulting from other coupling schemes are feasible taking into account the additivity of the CF effect with respect to 2^k -pole $\mathcal{H}_{CF}$ components. To achieve this one should employ consequently the tensor transformational properties of the states along with the standard angular momentum re-coupling techniques [3-5]. The systematic calculations have been performed in the paper for all the non-Kramers and Kramers states with $1 \leq J \leq 8$, it means for fifteen J values.

In order to compare the splitting effects of arbitrary (real or virtual) $\mathcal{H}_{CF}$ s they must yield the splitting diagrams of the same second moment σ^2 defined as [6-9]

$$\sigma^2 |J\rangle = \frac{1}{2J+1} \sum_n [E_n - \overline{E}(|J\rangle)]^2 = \frac{1}{2J+1} \sum_k S_k^2 \left(\langle J || C^{(k)} || J \rangle \right)^2, \quad (2)$$

where the energy centre of gravity of the sublevels within the state $|J\rangle$ is given by $\overline{E}(|J\rangle) = \frac{1}{2J+1} \sum_n E_n$, with E_n as the energy of $|n\rangle$ sublevel, and $S_k = \left(\frac{1}{2k+1} \sum_q |B_{kq}|^2 \right)^{1/2}$ is the conventional CF strength of the 2^k -pole [6,7,10,11], whereas the dimensionless scalar $\langle J || C^{(k)} || J \rangle$ describes the 2^k -pole type asphericity of the state $|J\rangle$ [2,9].

To put in order different isomodal 2^k -pole $\mathcal{H}_{CF}^{(k)}$ components of the global $\mathcal{H}_{CF}$ the complementary scale of the CF strength was applied. This is the spherically averaged modulus of $\mathcal{H}_{CF}^{(k)}$ which is equal to the spherically averaged modulus of the axial B_{k0} CFP [8]

$$|\mathcal{H}_{CF}^{(k)}|_{av} = \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi |\mathcal{H}_{CF}^{(k)}(\alpha, \beta)| \sin \beta d\beta d\alpha = |B_{k0}|_{av}, \quad (3)$$

where α and β are two Euler angles of the reference frame rotation (the third angle is inessential).

The most general $\mathcal{H}_{CF}$ contains 27 CFPs including 3 real axial parameters and 12 pairs of complex ones [2,12,13]. An interesting question arises whether one should consider complex parametrized $\mathcal{H}_{CF}$ s investigating the extreme splittings ΔE_{min} and ΔE_{max} of electron states in any CF yielding the same σ^2 . The negative answer can be explained as follows. The off-diagonal matrix elements of

the complex $\mathcal{H}_{\text{CFs}}$ are superpositions of at most three complex terms (for $k = 2, 4, 6$), which due to their own different phases can be treated as vector sums. On the other hand in the case of the real parametrizations we deal solely with algebraic sums of the components. In consequence, the complex parametrizations (with the phase angles as their additional degrees of freedom) considerably enlarge the set of available ratios between the matrix elements comparing to a more discrete set for the real parametrizations. For each complex parametrized $\mathcal{H}_{\text{CF}}$ an equivalent real CF interaction matrix, leading to the same splitting diagram, can be given. The point is, however, that such a matrix cannot be attained only by means of the real CFPs. The complex Hamiltonians are therefore more capable than the real ones. This is what a supremacy of the complex parametrizations over the real ones consists in. Since the absolute values of the vector sums are enclosed within the range of the absolute values of the relevant algebraic sums, $\mathcal{H}_{\text{CFs}}$ of complex parametrizations are not relevant to $\Delta E_{\min}$ and $\Delta E_{\max}$, although they essentially enrich the set of available splitting diagrams. Thus, searching for the extreme splittings we can confine ourselves exclusively to the $\mathcal{H}_{\text{CF}}$ of real CFPs. For the sake of simplicity the terms $\frac{1}{\sqrt{2}} (C_q^{(k)} + C_{-q}^{(k)})$ are written as $C_q^{(k)}$.

Clearly, the numerically-analytical approach applied previously to the three-parameter axial CFs, in this case seems to be hardly realizable considering the large number of independent CFPs. In fact, for fifteen real CFPs and fixed σ^2 one would need to scan independently and simultaneously all the involved B_{kq} s within the ranges from 0 to $\sqrt{(2J+1)(2k+1)}$ (in σ), and then, after diagonalization of the $\mathcal{H}_{\text{CF}}$ interaction matrix, to find out the upper and lower limits of the dominating absolute differences between the sublevels energies. In geometrical interpretation it would correspond to the analysis of adequate number of plane equations (equal to the number of the sublevels) within the 15-dimensional space with respect to the sphere of the radius $R = \sigma$ [1]. In order to avoid such difficulties we have investigated $\Delta E_{\min}$ and $\Delta E_{\max}$ for a general $\mathcal{H}_{\text{CF}}$ (Eq.(1)) employing an intermediate method. Namely, we utilized a seemingly trivial fact that the CF effect is essentially differentiated for various combinations of q and k indexes. Firstly, we have found the limits of the total splittings which can be obtained separately for isomodal partial $\mathcal{H}_{\text{CF}}^{(k)} = \sum_q B_{kq} C_q^{(k)}$ for $k = 2, 4$ and 6 (section 2), and next we have calculated the analogous limits for the potentials $\mathcal{H}_{\text{CF}}(q) = B_{2q} C_q^{(2)} + B_{4q} C_q^{(4)} + B_{6q} C_q^{(6)}$, this time with a fixed q index (section 3).

Collating the results from both the approaches we are able to deduce the ranges $(\Delta E_{\min}, \Delta E_{\max})$ binding all the crystal fields and the electron states $|J\rangle$ with $1 \leq J \leq 8$ (section 4). The information on the extreme total splittings is valuable and helpful for experimentalists attempting to assign or verify complicated energy level spectra in crystal-fields.

2. Dispersion of the total splitting ΔE of $|J\rangle$ electron states in CFs of individual $2^{(k)}$ -pole $\mathcal{H}_{\text{CF}}^{(k)}$ for $k = 2, 4, 6$ yielding the same σ^2 .

In order to find the limits of the observed total splittings ΔE and to compare the splitting capability of different isomodal $\mathcal{H}_{\text{CF}}^{(k)}$ with respect to $|J\rangle$ states the relevant interaction matrices of order $2J+1$ for all $1 \leq J \leq 8$ have been diagonalized. There were chosen 8, 25 and 30 various $\mathcal{H}_{\text{CF}}^{(k)}$ s, both real and virtual, for $k = 2, 4$ and 6 , respectively.

These intentionally selected representative potentials are characterized by different contributions of the particular q -components. The exact and detailed quantitative compositions of the considered potentials are comprised in Appendix. The calculated minimal and maximal total splittings, their averages and mean square deviations are compiled in Tables 1, 2 and 3, respectively for $k = 2, 4$ and 6 . The complementary CF strength scale [8], i.e. the spherically averaged absolute values of $|\mathcal{H}_{\text{CF}}^{(k)}|_{\text{av}}$, enabled us to list in a systematic way the different isomodal $\mathcal{H}_{\text{CF}}^{(k)}$ s. However, its correlation with ΔE is not monotonic and direct.

Our initial assumption comparing the splitting effects of various $\mathcal{H}_{\text{CF}}^{(k)}$ s is the constant second moment σ^2 of the observed splittings instead of the relevant conventional CF strength [6,7,10,11]. Nevertheless, considering the $\mathcal{H}_{\text{CF}}^{(k)}$ compositions the latter invariant is more convenient. The energy quantities such as B_{kq} , ΔE , $|\mathcal{H}_{\text{CF}}^{(k)}|_{\text{av}}$, and δ take in these two approaches different values but they are correlated by a simple linear scaling relationship $\sigma_k^2 = S_k^2 \frac{(\langle J||C^{(k)}||J \rangle)^2}{(2J+1)(2k+1)}$ [6-9]. Further, for the sake of simplicity, writing the CF Hamiltonians we assume that $S_k^2 = 1$ and ignore the scalar factor $\frac{\sqrt{(2J+1)(2k+1)}}{\langle J||C^{(k)}||J \rangle}$. The last one ought to be introduced to ensure the condition $\sigma_k^2 = 1$.

However, even the most careful and attentive choice of representative isomodal $\mathcal{H}_{\text{CF}}^{(k)}$ sets covering the entire range of known calculated $|\mathcal{H}_{\text{CF}}^{(k)}|_{\text{av}}$ and preceded by several surveys, does not guarantee, in general case, to find the exact extreme splittings $\Delta E_{\min}$ and $\Delta E_{\max}$. The extreme total splittings proved to be certain have been found only for the quadrupolar $\mathcal{H}_{\text{CF}}^{(2)}$ acting on all the $|J\rangle$ states, as well as in some particular cases for the higher CF multipoles (Tables 1-3 vs. Table 10). Since the infinite number of $\mathcal{H}_{\text{CF}}^{(k)}$ s cannot be explicitly inspected, the minimal and maximal total splittings found numerically for the considered $\mathcal{H}_{\text{CF}}^{(k)}$ sets have been assumed as the extreme ones. Thus, in most cases they have a hypothetical character of statistical origin due to their sample nature. Fortunately, this intra-multipolar (for a fixed k) q -mixing mechanism does not determine the extreme total splittings of the global $\mathcal{H}_{\text{CF}} = \sum_k \mathcal{H}_{\text{CF}}^{(k)}$ (section 3 and 4). The exact numerical approach consisting in mapping of all the independent CFPs variability ranges would be a generalization of the previous technique employed for the axial CFs [1]. It is however a hardly realizable task due to its dimensionality even for an individual k . Our more practical choice to overcome this complexity was a numerical experiment that does not guarantee the absolute character of the found extreme values, although there are some convincing premises making them reliable. The apparent over-representation of the simple mono-

parametric $\mathcal{H}_{\text{CF}}^{(k)}$ in Tables 1-3 leads to the conclusion that the postulated $\Delta E_{\min}$ and $\Delta E_{\max}$ should be correct or at least close to the actual extreme values. Still better consistence should hold for the ΔE averages and the mean square deviations δ . The CFP mixing within the individual 2^k -poles, i.e. within the component $\mathcal{H}_{\text{CF}}^{(k)}$, generally does not promote the extreme ΔE . The reason for that is the additivity of σ^2 with respect to $|B_{kq}|^2$ (Eq. (2)). It means that every B_{kq} contributes separately and independently to σ^2 producing however qualitatively various partial splittings. Considering the linear independence of the involved operators $C_q^{(k)}$ for different q , which implies from their orthogonality, the resultant CF effect should come rather from the average of the partial contributions. However, it does not exclude some particular superpositions corresponding e.g. to high-symmetrical $\mathcal{H}_{\text{CF}}^{(k)}$ s, such as cubic potentials, which promote rather symmetric dichotomic splittings with ΔE approaching 2σ . In order to distinguish the statistical data compiled in Tables 1-3 from the exact results obtained strictly numerically (section 3), the former are given with the limited accuracy of $10^{-2}\sigma$ whereas the $\overline{\Delta E}$ and δ with the accuracy of $10^{-3}\sigma$. A review of the chosen $\mathcal{H}_{\text{CF}}^{(k)}$ s with respect to the produced extreme splittings is discussed below.

In the case of the pure quadrupolar CFs, $\mathcal{H}_{\text{CF}}^{(2)}$, (Table 1 and Appendix) for all the $|J\rangle$ states with $1 \leq J \leq 8$, the $\Delta E_{\min}$ is attained under the action of the $\mathcal{H}_{\text{CF}}^{(2)} = C_0^{(2)}$ Hamiltonian (further denoted by the $[C_0^{(2)}]\text{CF}$), whereas $\Delta E_{\max}$ either in $[C_2^{(2)}]$ or $[C_1^{(2)}]\text{CF}$ s. Only for $\mathcal{H}_{\text{CF}}^{(2)}$ s the unambiguous and monotonic relationship between ΔE and the relevant $\overline{|E_n|}$ is observed (Appendix). The $\Delta E_{\min}$ corresponds to the largest $\overline{|E_n|}$, whereas the $\Delta E_{\max}$ to the smallest $\overline{|E_n|}$ (what is understood for the $\sigma^2 = \text{const}$ condition), and these magnitudes are the nominally extreme ones, indeed. In the quadrupolar case the mean $\overline{\Delta E}$ for the chosen eight representative $\mathcal{H}_{\text{CF}}^{(2)}$ s rises monotonically from 2.343σ for $J = 1$ up to 3.185σ for $J = 8$ (except the trivial case of $J = 3/2$ when it is always 2.0000σ) and is shifted towards the $\Delta E_{\max}$. The mean square deviation δ is of the order of 0.1σ and becomes much smaller within the range $3/2 \leq J \leq 4$ (Table 1). For $J = 3/2$ and 2 the ΔE is entirely independent of the $\mathcal{H}_{\text{CF}}^{(2)}$ composition, whereas for $J = 5/2$ and 3 depends only slightly with the δ being merely of the order of 0.01σ .

In the case of the 2^4 -pole, $\mathcal{H}_{\text{CF}}^{(4)}$, (Table 2 and Appendix) for the noticeable part of J values ($J = 2, 5/2, 11/2 \div 8$) the $\Delta E_{\min}$ is achieved in the cubic CF: $[\frac{1}{2}\sqrt{\frac{7}{3}}C_0^{(4)} + \frac{1}{2}\sqrt{\frac{5}{3}}C_4^{(4)}]$ confirming its tendency to the dichotomic type splittings. In turn, for $J = 3$ and $9/2$ the $\Delta E_{\min}$ takes place in $[C_4^{(4)}]\text{CF}$, and for $J = 7/2$ and 4 in $[\frac{1}{\sqrt{2}}C_0^{(4)} + \frac{1}{\sqrt{2}}C_2^{(4)}]\text{CF}$, although the approximate magnitudes for both the minima occur in $[C_4^{(4)}]\text{CF}$. The case of $J = 5$ seems to be quite intriguing since the ΔE is completely independent on the $\mathcal{H}_{\text{CF}}^{(4)}$ composition what means here that $\delta = 0$. Further, the $\Delta E_{\max}$ for $J = 2$ and $5/2$ is gained in $[C_4^{(4)}]\text{CF}$, for $J = 3 \div 9/2$ in $[C_3^{(4)}]\text{CF}$, for $J = 11/2 \div 13/2$ in $[C_0^{(4)}]\text{axial CF}$, next for $J = 7$ in $[\frac{1}{\sqrt{2}}C_2^{(4)} + \frac{1}{\sqrt{2}}C_3^{(4)}]\text{CF}$, and finally for $J = 15/2$ and 8 in $[C_1^{(4)}]\text{CF}$. It turns out that the same CF, $[C_0^{(4)}]$, yields $\Delta E_{\min}$ for the states with $J = 3 \div 9/2$, as well as the $\Delta E_{\max}$ for the states with $J = 2$ and $5/2$. It is easy to show on the splitting diagrams and taking into account

the relevant crystal-quantum numbers. The mean $\overline{\Delta E}$ magnitudes do not change considerably with J oscillating around 2.60σ . What is worthy to notice is the constant $\Delta E = 2.3531\sigma$ for $J = 5$ and dominating $\overline{\Delta E} = 2.965\sigma$ for $J = 3$. In the case of $k = 4$ the mean square deviation δ is generally somewhat higher than for $k = 2$, but also of the order of 0.1σ and rises apparently for $J = 2$ and 3 decreasing within the range $9/2 \leq J \leq 6$.

In the case of the 2^6 -pole, $\mathcal{H}_{\text{CF}}^{(6)}$, (Table 3 and Appendix) for $J = 3$ and 4 the $\Delta E_{\min}$ has been found in $[C_3^{(6)}]\text{CF}$, for $J = 7/2$ and 5 in the cubic CF: $[\frac{1}{2\sqrt{2}}C_0^{(6)} - \frac{\sqrt{7}}{2\sqrt{2}}C_4^{(6)}]$, but in turn for $J = 9/2$ in $[\frac{1}{\sqrt{13}}C_0^{(6)} + \sqrt{\frac{2}{13}}C_1^{(6)} + \sqrt{\frac{2}{13}}C_2^{(6)} + \sqrt{\frac{2}{13}}C_3^{(6)} - \sqrt{\frac{2}{13}}C_4^{(6)} + \sqrt{\frac{2}{13}}C_5^{(6)} + \sqrt{\frac{2}{13}}C_6^{(6)}]\text{CF}$ of rather uniform contribution of all the q -components. Its effect is however close to that in $[C_3^{(6)}]\text{CF}$. Finally, for $J = 11/2 \div 8$ the $\Delta E_{\min}$ is gained in $[C_6^{(6)}]\text{CF}$. In turn, the $\Delta E_{\max}$ for $J = 3$ and $7/2$ is attained in $[C_6^{(6)}]\text{CF}$, and for $J = 4$ in $[C_4^{(6)}]\text{CF}$. For $J = 9/2$ a rather isolated $\Delta E_{\max}$ has been found in $[\frac{1}{\sqrt{3}}C_1^{(6)} + \frac{1}{\sqrt{3}}C_3^{(6)} + \frac{1}{\sqrt{3}}C_5^{(6)}]\text{CF}$. For $J = 5$ it occurs in $[C_2^{(6)}]\text{CF}$, but its effect is almost identical with that in $[C_3^{(6)}]$ or $[C_5^{(6)}]\text{CFs}$. In turn, in the case of $J = 6$ it appears in $[\frac{1}{\sqrt{13}}C_0^{(6)} + \sqrt{\frac{2}{13}}C_1^{(6)} + \sqrt{\frac{2}{13}}C_2^{(6)} + \sqrt{\frac{2}{13}}C_3^{(6)} + \sqrt{\frac{2}{13}}C_4^{(6)} + \sqrt{\frac{2}{13}}C_5^{(6)} - \sqrt{\frac{2}{13}}C_6^{(6)}]\text{CF}$. At last, for $J = 7$ and $15/2$ the $\Delta E_{\max}$ occurs in $[\frac{1}{\sqrt{2}}C_0^{(6)} + \frac{1}{\sqrt{2}}C_6^{(6)}]\text{CF}$, and finally in the case of $J = 8$ it appears in $[\frac{1}{\sqrt{13}}C_0^{(6)} - \sqrt{\frac{2}{13}}C_1^{(6)} + \sqrt{\frac{2}{13}}C_2^{(6)} + \sqrt{\frac{2}{13}}C_3^{(6)} + \sqrt{\frac{2}{13}}C_4^{(6)} + \sqrt{\frac{2}{13}}C_5^{(6)} + \sqrt{\frac{2}{13}}C_6^{(6)}]\text{CF}$, which is close to the ΔE in $[C_2^{(6)}]\text{CF}$. The mean value of ΔE slightly changes with J oscillating around 2.95σ and is somewhat enhanced for $J = 4$ and 6 . The mean square deviation fluctuates around 0.2σ reaching its maximal value $\delta = 0.36\sigma$ for $J = 3$, and the minimal value $\delta = 0.12\sigma$ for $J = 9/2$.

Taking into account the fundamental differences between the three effective 2^k -poles contributing to the global $\mathcal{H}_{\text{CF}}$ the observed discrepancies in ΔE , $\overline{\Delta E}$, and δ for the particular multipoles are surprisingly small with respect to both their magnitudes as well as J dependence (Tables 1 - 3). Apart from the appreciable increase in δ roughly in ratio $1 : 2 : 3$ for $k = 2, 4$ and 6 , respectively, all the three additive contributions can be, in principle, considered as comparable. The greater dispersion of the ΔE (it means the greater δ) for higher k corresponds to the higher filling of the $(2J + 1)$ -dimensional interaction matrices by the non-zero $\mathcal{H}_{\text{CF}}$ matrix elements.

The observed ΔE fluctuations result primarily from the arithmetic relations (stemming from the quantization consequences) between the involved numbers M_J , $|M_J| \leq J$, and q , $|q| \leq k$, according to the definition of the crystal quantum number μ , $M_J = \mu(\text{mod } q)$ [2,14]. For example, the state $|J = 3\rangle$ in $[C_6^{(6)}]\text{CF}$ becomes split into two singlets ($\mu = 3$ and 3), whereas all the other substates $|M_J\rangle$ remain intact. In consequence, the maximal nominally admissible total splitting $\Delta E = \Delta\mathcal{E} = 3.7417\sigma$ is attained (Table 10). However, exactly the same potential splits the state $|J = 6\rangle$ into four doublets ($\mu = 1, 2, 4$ and 5) and five singlets ($\mu = 0, 0, 0, 3$ and 3) acting on all the initial substates what results in the low value of $\Delta E = 2.6000\sigma$ – the lowest one found for the state, exceeding however the nominally minimal $\Delta\mathcal{E} = 2.0059\sigma$. A comparison of the extreme and average total splittings given in Tables 1 - 3 along with the relevant axial splittings for $k = 2, 4$ and 6 presented on the left side of

Table 4 is instructive because it demonstrates the position of the $\mathcal{H}_{\text{CF}}^{(k)} = C_0^{(k)}$ among the other $\mathcal{H}_{\text{CF}}^{(k)}$ potentials.

Above we have investigated the effect of the $|J\rangle$ states splitting by the three individual 2^k -poles separately. Nevertheless, under the action of the global $\mathcal{H}_{\text{CF}} = \sum_{k=2,4,6} \mathcal{H}_{\text{CF}}^{(k)}$, yielding the same σ^2 , there appears the second mechanism effectively leading to the dispersion of the available ΔE . This is the resultant effect of the $C_q^{(k)}$ components with the same q but different k (three components for $q = 0, 1$ and 2, and two ones for $q = 3$ and 4). And this is the mechanism which turns out to be decisive. Its thorough analysis is consistently provided in the next section.

3. Dispersion of the total splitting ΔE of $|J\rangle$ states in CFs with fixed q : $\mathcal{H}_{\text{CF}}(q) = B_{2q}C_q^{(2)} + B_{4q}C_q^{(4)} + B_{6q}C_q^{(6)}$ yielding the same σ^2 .

For all the $\mathcal{H}_{\text{CF}}^{(k)}$ s the splitting capability of the mono-parametric Hamiltonians (with fixed $|q|$) seems to be dominating or almost dominating. Therefore, in our the next step of the search for the extreme total splittings we have focused on the q -component superpositions $\mathcal{H}_{\text{CF}}(q) = B_{2q}C_q^{(2)} + B_{4q}C_q^{(4)} + B_{6q}C_q^{(6)}$ with fixed q running from 0 to 4. They have been optimized with respect to their splitting capability. Obviously, the assumption that the $\mathcal{H}_{\text{CF}}(q)$ is normalized, leading to the same constant σ^2 , still remains in power.

In the case of a three-parameter $\mathcal{H}_{\text{CF}}(q)$ form, the method applied previously to the axial CFs [1] can be adapted here as well. However, because of the non-diagonality of the $\mathcal{H}_{\text{CF}}(q)$ interaction matrices for $q \neq 0$, the mapping procedure has to be preceded by their diagonalization. The splitting effect of each 2^k -pole superposition has been analyzed within the three-dimensional spherical reference frame (R, θ, φ) . Taking the radius of the sphere $R = \sigma$, the matrix elements have the form (in σ):

$$\begin{aligned} \langle JM_J | \mathcal{H}_{\text{CF}}(q) | JM'_J \rangle &= (-1)^{J-M_J} \sqrt{5(2J+1)} \begin{pmatrix} J & 2 & J \\ M_J & q & M'_J \end{pmatrix} \sin \theta \cos \varphi \\ &+ (-1)^{J-M_J} 3\sqrt{2J+1} \begin{pmatrix} J & 4 & J \\ M_J & q & M'_J \end{pmatrix} \sin \theta \sin \varphi \\ &+ (-1)^{J-M_J} \sqrt{13(2J+1)} \begin{pmatrix} J & 6 & J \\ M_J & q & M'_J \end{pmatrix} \cos \theta, \end{aligned} \quad (4)$$

where the coordinates $0 \leq \theta < \pi$ and $0 \leq \varphi < 2\pi$ define the multipolar composition of the $\mathcal{H}_{\text{CF}}(q)$. In the case of the two-parameter superpositions for $q = 3$ or 4 the above formula reduces itself to the simplified form which can be obtained substituting $\varphi = \pi/2$ into Eq.(4) (due to the lack of the quadrupolar term).

After diagonalization of the matrices, i.e. having the sublevels eigenvalues, and mapping the whole variation ranges of the coordinates θ and φ , the upper and lower limits of the dominating

absolute differences among the eigenvalues at each (θ, φ) point have been calculated numerically. In consequence, the physically admissible total splittings ΔE_{min} and ΔE_{max} of the $|J\rangle$ states in CFs yielding the same σ^2 have been found. The whole ranges of the θ and φ angles digitized with the accuracy of $2 \cdot 10^{-4}$ and $1 \cdot 10^{-4}$, respectively, has been swept up numerically. The evaluation of the upper and lower limits has been facilitated by a short Fortran programme [1].

The calculated ΔE_{min} and ΔE_{max} for $q = 0 \div 4$ are enclosed in Tables 4 - 8, respectively. To complete these data and make them easy to compare the relevant ΔE for the solitary contributions to $\mathcal{H}_{CF}$ (i.e. for $q = 5$ or 6) are appended in Table 9.

The essential results stand as follows:

- There are three types of the $\mathcal{H}_{CF}(q)$ with fixed q : the first involving all the three terms with $k = 2, 4$ and 6 for $q = 0, 1$ and 2 , the second composed of two terms with $k = 4$ and 6 for $q = 3$ and 4 , and the third based only on the one term with $k = 6$ for $q = 5$ and 6 . Intuitively, the three-multipolar $\mathcal{H}_{CF}$ s are the most effective in achieving the extreme total splittings. The results confirm it convincingly. From among all the 30 extreme values merely 5 (three ΔE_{max} s and two ΔE_{min} s) come from the two-multipolar $\mathcal{H}_{CF}$ s (three for $\mathcal{H}_{CF}(q = 4)$ and two for $\mathcal{H}_{CF}(q = 3)$) (see Tables 4 - 9). From among the complete three-multipolar superpositions only $\mathcal{H}_{CF}(q = 1)$ s lead for $1 \leq J \leq 8$ (except for $J = 4$) to both the largest ΔE_{max} and the largest ΔE_{min} . In turn, the three-multipolar $\mathcal{H}_{CF}(q = 0)$ s lead to the smallest ΔE_{min} as well as to the smallest ΔE_{max} (again except for $J = 4$). It confirms unambiguously the extreme splitting capabilities of these two particular CF potentials.
- There exist $\mathcal{H}_{CF}(q = 1)$ s for which the ΔE_{max} for all the considered Kramers states with $J \leq 15/2$ practically reach the maximal, nominally predicted values (compare Table 5 vs. Table 10). The observed deviation found for $J = 15/2$ amounts to 0.3% ($\Delta E_{max} = 3.9880\sigma$ vs. $\Delta \mathcal{E}_{max} = 4.0000\sigma$), and for $J = 13/2$ only 0.1% ($\Delta E_{max} = 3.7381\sigma$ vs. $\Delta \mathcal{E}_{max} = 3.7417\sigma$). These discrepancies follow the characteristic divergence between the ΔE_{max} and $\Delta \mathcal{E}_{max}$ values appearing for large J values due to breaking of the simple model relation $\Delta E_{max} = \sigma\sqrt{2J+1}$. It means that the splitting diagrams with two doublets at energies $\pm\Delta E/2$ and all the others at zero energy become unavailable. For the $\mathcal{H}_{CF}(q = 2)$ s and the half-integer J the dependence $\Delta E_{max}(J)$ approximates closely that for the $\mathcal{H}_{CF}(q = 1)$ s discussed above (Tables 5 and 6).
- For the non-Kramers states the ΔE_{max} is also reached for the majority of J numbers under the action of the $\mathcal{H}_{CF}(q = 1)$ s. Here, only three exceptions have been found. For $J = 4$ the prevailing $\Delta E_{max} = 4.0763\sigma$ occurs for the $\mathcal{H}_{CF}(q = 4)$ (Table 8), and in the case of $J = 5$ and 6 for the $\mathcal{H}_{CF}(q = 3)$ s, where ΔE_{max} attains 3.5904σ and 3.7054σ , respectively. The latter two values only slightly exceed their counterparts for the $\mathcal{H}_{CF}(q = 1)$ s (Tables 7 and 5).

- The smallest total splittings ΔE_{min} both for the Kramers and non-Kramers ions have been found in the axial CFs for the $\mathcal{H}_{CF}(q=0)$ (Table 4). Only for $J = 15/2$ and 8 somewhat lower ΔE_{min} have been achieved for the $\mathcal{H}_{CF}(q=4)$ (Table 8).

4. Discussion

The key issue for the estimation of the maximal splittings of electron states in CFs is the fact that the maximal nominally predicted $\Delta \mathcal{E}_{max}$ for all the Kramers states with $J \leq 15/2$ are attained for the properly composed multipolar superpositions in $\mathcal{H}_{CF}(q=1) = B_{21}C_1^{(2)} + B_{41}C_1^{(4)} + B_{61}C_1^{(6)}$ (compare Tables 5 and 10). For the Kramers states the nominally allowed extreme $\Delta \mathcal{E}$ (resulting solely from the σ^2 constancy requirement [1,8]) amount to: $\Delta \mathcal{E}_{max} = \sigma\sqrt{2J+1}$ and $\Delta \mathcal{E}_{min} = 2\sigma$ for $J = 3/2, 7/2, 11/2$ and $15/2$, and $\Delta \mathcal{E}_{min} = \sigma \frac{2(2J+1)}{\sqrt{(2J+3)(2J-1)}}$ for $J = 5/2, 9/2$ and $13/2$. In turn, for the non-Kramers states they are: $\Delta \mathcal{E}_{max} = \sigma\sqrt{2(2J+1)}$ and $\Delta \mathcal{E}_{min} = \sigma \frac{2J+1}{\sqrt{J(J+1)}}$ (see Table 10).

It means, that no other $\mathcal{H}_{CF}$, of any unrestricted composition, can produce larger splittings. This fact entitles us to recognize the $\mathcal{H}_{CF}(q=1)$ as the strongest potential with respect to the produced total splitting from among all the possible $\mathcal{H}_{CF}$ s yielding the same σ^2 . It confirms also the secondary role of the q -mixing mechanism of the $C_q^{(k)}$ components in the $\mathcal{H}_{CF}^{(k)}$ (Tables 1-3), as well as the q and k cross-mixing mechanism in the $\mathcal{H}_{CF}$.

Similarly, for the non-Kramers states the $\mathcal{H}_{CF}(q=1)$ ensures the dominating ΔE_{max} (Table 5), and the found exceptions for $J = 4 \div 6$ and $q = 3$ and 4 result presumably from more favourable configurations of the sublevels for the involved crystal-quantum numbers (section 2). What promotes the large ΔE_{max} is putting two of the sublevels (best of all two singlets) possibly far away from the energy centre of gravity and bringing together the remaining ones in the vicinity of the centre.

The $\mathcal{H}_{CF}(q=1)$ as the only one does not allow doublets to occur in the energy spectrum (ignoring an accidental degeneracy) and splits the $|J\rangle$ state into $2J+1$ singlets. In contrary, the $\mathcal{H}_{CF}(q=0)$ splits the $|J\rangle$ state always into J doublets and only one singlet $|M_J=0\rangle$. The $\mathcal{H}_{CF}(q)$ s for $q = 2 \div 6$ generate mixtures of doublets and singlets. The splitting diagrams for the $\mathcal{H}_{CF}(q \neq 1)$ s are symmetrical with respect to their centres of gravity, which means that the singlet $|M_J=0\rangle$ in the non-Kramers states has to always lie at the centre of gravity. This symmetry does not hold for $q=0$.

In the light of the above results we postulate that the ΔE_{max} found for the $\mathcal{H}_{CF}(q=1)$ s completed by the dominating values for the $\mathcal{H}_{CF}(q=4)$ in the case of $J=4$, as well as for the $\mathcal{H}_{CF}(q=3)$ in the case of $J=5$ and 6, are the highest limits of the total CF splittings of the $|J\rangle$ states for $1 \leq J \leq 8$ (Table 11).

The lower limits of the ΔE_{min} are achievable presumably in the axial CFs of the $\mathcal{H}_{CF}(q=0)$. Firstly, these are the smallest values from all the presented in Tables 4 - 9, except slightly smaller ΔE_{min} s for $J = 15/2$ and 8 under the action of $\mathcal{H}_{CF}(q=4)$. Secondly, the axial CFs split the

non-Kramers states into the highest number of doublets. This qualitative similarity of the splitting diagrams of the non-Kramers and Kramers states in the axial CFs must result in the equally small extreme total splittings (Table 4). Moreover, the obtained ΔE_{min} s do not differ distinctly (contrary to the ΔE_{max} s for higher J) from their nominal limits (Table 10), although the energy distances should not be treated linearly. The particular weight of the axial CFs results also from their leading role in the complementary scale of the CF strength [8]. For all the three 2^k -poles the $|\mathcal{H}_{CF}^{(k)}|_{av}$ values in the axial CFs are the largest in the sets of all the remaining CFs (Appendix). This suggests rather higher average magnitudes of the absolute energies $\overline{E_n}$ of the sublevels and naturally lower ΔE .

Thus, we postulate that the ΔE_{min} achieved in the axial CFs of the $\mathcal{H}_{CF}(q = 0)$ for $1 \leq q \leq 7$ and in the CFs of the $\mathcal{H}_{CF}(q = 4)$ for $J = 15/2$ and 8 are the lowest possible limits of the total CF splittings of the $|J\rangle$ states. All the above estimated maximal and minimal total splittings of the $|J\rangle$ states for $1 \leq q \leq 8$ in any CF are compiled in Table 11.

Concluding, two types of the CF Hamiltonians can be distinguished with respect to their extreme splitting capability. The lower limits of the total splittings are attained in the prevailing number of cases in the axial CFs of the $\mathcal{H}_{CF}(q = 0) = B_{20}C_0^{(2)} + B_{40}C_0^{(4)} + B_{60}C_0^{(6)}$, whereas the upper limits in the CFs of the $\mathcal{H}_{CF}(q = 1) = B_{21}C_1^{(2)} + B_{41}C_1^{(4)} + B_{61}C_1^{(6)}$. For the non-Kramers states in the first case we deal with the maximal number of doublets (equal to J), whereas in the second case with the maximal number of singlets (equal to $2J + 1$). In turn, for the Kramers states only doublets occur. These two particular Hamiltonians composed of the $C_q^{(k)}$ operators with the same q (either 0 or 1) turn out to be the most effective from the extreme total splittings viewpoint. They contribute to the same elements of the $\mathcal{H}_{CF}(q)$ interaction matrices. On the other hand, the components $C_q^{(k)}$ with different q contribute to various matrix elements. This leads to more advanced mixing of the initial free ion substates which averages the angle distributions of their electron density. Such effect does not however favour the extreme splittings.

The lower limits of the total splittings ΔE_{min} lie within the range $(2.0000\sigma, 2.4081\sigma)$, whereas the nominally allowed $\Delta \mathcal{E}_{min}$ in $(2.0000\sigma, 2.1213\sigma)$.

The upper limits ΔE_{max} can change within the range $(2.0000\sigma, 4.1043\sigma)$ compared with the nominally allowed $\Delta \mathcal{E}_{max}$ interval $(2.0000\sigma, 5.8310\sigma)$. The gap $\Delta E_{max} - \Delta E_{min}$ varies from 0 to 1.9961 (for $J = 4$) and is smaller for the Kramers states (Table 11).

It is worthy to remind that all the energy quantities met in the paper are referred to constant σ^2 and expressed in σ . However, the σ depends not only on the CFPs (Eq.(2)) but in equal degree on the multipolar characteristics of the electron density distribution of the central ion affected by the CF.

References

- [1] J. Mulak, M. Mulak, Phys. Status Solidi (b) 1-8 (2009), DOI 10.1002/pssb.200945087.
- [2] B. G. Wybourne, Spectroscopic Properties of Rare Earths (John Wiley, New York, 1965).
- [3] B. R. Judd, Operator Techniques in Atomic Spectroscopy (Mc Graw-Hill, New York, 1963).
- [4] A. R. Edmonds, Angular Momentum in Quantum Mechanics (Princeton University Press, Princeton, New York, 1960).
- [5] M. Rotenberg, R. Bivins, N. Metropolis and J.K. Wooten, Jr., The 3-j and 6-j Symbols (MIT Press, Cambridge, MA, 1963).
- [6] Y. Y. Yeung, Invariants and moments in: Crystal Field Handbook, edited by D.J. Newman and B. Ng (Cambridge University Press, Cambridge, MA, 2000).
- [7] R. P. Leavitt, J. Chem Phys. **77**, 1661 (1982).
- [8] J. Mulak, M. Mulak, J. Phys. A: Math. Theor. **40**, 1 (2007).
- [9] J. Mulak, M. Mulak, Phys. Stat. Solidi B **245**, 1156 (2008).
- [10] F. Auzel and O.L. Malta, J. Phys. (France) **44**, 201 (1983).
- [11] C. Rudowicz and J. Qin, J. Lumin. **110**, 39 (2004).
- [12] C. Görller-Walrand and K. Binnemans, Rationalization of crystal-field parametrization, in: Handbook of the Physics and Chemistry of Rare Earths, edited by K.A. Gschneider, Jr. and L. Eyring (Elsevier, Amsterdam, 1996), Vol. 23.
- [13] J. Mulak and Z. Gajek, The Effective Crystal-Field Potential (Elsevier, Amsterdam, 2000).
- [14] K. H. Hellwege, Annalen der Physik, **4**, 95 (1949).

Appendix

Normalized ($S_k^2 = 1$) values of CFPs in the considered $\mathcal{H}_{\text{CF}}^{(k)}$ for $k = 2, 4$ and 6 given under the scheme $(B_{k0}, B_{k1}, B_{k-1}, B_{k2}, B_{k-2}, \dots, B_{kk}, B_{k-k})$ and the relevant total splitting ΔE and average absolute value of E_n , $|\overline{E_n}|$, for the $|J\rangle$ states given under the scheme $\mathbf{J}(\Delta E, |\overline{E_n}|)$. The Hamiltonians are ordered according to their complementary CF strength $|\mathcal{H}_{\text{CF}}^{(k)}|_{\text{av}}$ [8].

No.	$\mathcal{H}_{\text{CF}}^{(2)}$	$ \mathcal{H}_{\text{CF}}^{(2)} _{\text{av}}$
1	$(1, 0, 0, 0, 0)$ $\mathbf{1}(2.1213, 0.9427)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.9560)$, $\mathbf{5/2}(2.4056, 0.8911)$ $\mathbf{3}(2.5983, 0.8247)$, $\mathbf{7/2}(2.6184, 0.8728)$, $\mathbf{4}(2.7356, 0.8862)$, $\mathbf{9/2}(2.7540, 0.8811)$ $\mathbf{5}(2.8308, 0.8647)$, $\mathbf{11/2}(2.8443, 0.8536)$, $\mathbf{6}(2.9008, 0.8675)$, $\mathbf{13/2}(2.9124, 0.8718)$ $\mathbf{7}(2.9549, 0.8686)$, $\mathbf{15/2}(2.9641, 0.8604)$, $\mathbf{8}(2.9973, 0.8593)$.	axial 0.385
2	$(\frac{1}{2}, 0, 0, \frac{\sqrt{6}}{4}, \frac{\sqrt{6}}{4})$ $\mathbf{1}(2.1213, 0.9427)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.9560)$, $\mathbf{5/2}(2.4056, 0.8911)$ $\mathbf{3}(2.5983, 0.8247)$, $\mathbf{7/2}(2.6184, 0.8728)$, $\mathbf{4}(2.7356, 0.8862)$, $\mathbf{9/2}(2.7540, 0.8811)$ $\mathbf{5}(2.8308, 0.8647)$, $\mathbf{11/2}(2.8443, 0.8536)$, $\mathbf{6}(2.9008, 0.8675)$, $\mathbf{13/2}(2.9124, 0.8718)$ $\mathbf{7}(2.9549, 0.8686)$, $\mathbf{15/2}(2.9641, 0.8604)$, $\mathbf{8}(2.9973, 0.8593)$.	0.385
3	$(\frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, -\frac{1}{\sqrt{5}}, -\frac{1}{\sqrt{5}})$ $\mathbf{1}(2.3172, 0.9253)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.9475)$, $\mathbf{5/2}(2.4198, 0.8796)$ $\mathbf{3}(2.6090, 0.8283)$, $\mathbf{7/2}(2.6633, 0.8696)$, $\mathbf{4}(2.7759, 0.8794)$, $\mathbf{9/2}(2.8284, 0.8719)$ $\mathbf{5}(2.9012, 0.8543)$, $\mathbf{11/2}(2.9481, 0.8497)$, $\mathbf{6}(2.9992, 0.8619)$, $\mathbf{13/2}(3.0387, 0.8643)$ $\mathbf{7}(3.0770, 0.8591)$, $\mathbf{15/2}(3.1081, 0.8506)$, $\mathbf{8}(3.1383, 0.8540)$.	0.381
4	$(\frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}})$ $\mathbf{1}(2.4156, 0.8834)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.9260)$, $\mathbf{5/2}(2.4401, 0.8544)$ $\mathbf{3}(2.6256, 0.8830)$, $\mathbf{7/2}(2.7158, 0.8652)$, $\mathbf{4}(2.8255, 0.8647)$, $\mathbf{9/2}(2.9027, 0.8521)$ $\mathbf{5}(2.9769, 0.8351)$, $\mathbf{11/2}(3.0364, 0.8435)$, $\mathbf{6}(3.0903, 0.8490)$, $\mathbf{13/2}(3.1358, 0.8467)$ $\mathbf{7}(3.1774, 0.8392)$, $\mathbf{15/2}(3.2137, 0.8363)$, $\mathbf{8}(3.2462, 0.8427)$.	0.374
5	$(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}})$ $\mathbf{1}(2.4260, 0.8737)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.9210)$, $\mathbf{5/2}(2.4428, 0.8490)$ $\mathbf{3}(2.6273, 0.8336)$, $\mathbf{7/2}(2.7221, 0.8646)$, $\mathbf{4}(2.8322, 0.8613)$, $\mathbf{9/2}(2.9112, 0.8478)$ $\mathbf{5}(2.9850, 0.8343)$, $\mathbf{11/2}(3.0465, 0.8428)$, $\mathbf{6}(3.0999, 0.8465)$, $\mathbf{13/2}(3.1467, 0.8434)$ $\mathbf{7}(3.1878, 0.8348)$, $\mathbf{15/2}(3.2244, 0.8354)$, $\mathbf{8}(3.2573, 0.8399)$.	0.373
6	$(\frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}e^{i\pi/4}, \frac{1}{\sqrt{5}}e^{-i\pi/4}, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}})$ $\mathbf{1}(2.4450, 0.8431)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.9055)$, $\mathbf{5/2}(2.4483, 0.8314)$ $\mathbf{3}(2.6321, 0.8354)$, $\mathbf{7/2}(2.7341, 0.8627)$, $\mathbf{4}(2.8436, 0.8519)$, $\mathbf{9/2}(2.9267, 0.8344)$ $\mathbf{5}(3.0013, 0.8314)$, $\mathbf{11/2}(3.0643, 0.8404)$, $\mathbf{6}(3.1193, 0.8385)$, $\mathbf{13/2}(3.1668, 0.8325)$ $\mathbf{7}(3.2086, 0.8288)$, $\mathbf{15/2}(3.2468, 0.8327)$, $\mathbf{8}(3.2794, 0.8334)$.	0.369
7	$(0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0)$ $\mathbf{1}(2.4495, 0.8164)$, $\mathbf{3/2}(2.0000, 1.0000)$, $\mathbf{2}(2.3905, 0.8920)$, $\mathbf{5/2}(2.4495, 0.8164)$ $\mathbf{3}(2.6332, 0.8354)$, $\mathbf{7/2}(2.7373, 0.8627)$, $\mathbf{4}(2.8470, 0.8439)$, $\mathbf{9/2}(2.9310, 0.8238)$ $\mathbf{5}(3.0058, 0.8299)$, $\mathbf{11/2}(3.0682, 0.8404)$, $\mathbf{6}(3.1233, 0.8328)$, $\mathbf{13/2}(3.1709, 0.8241)$ $\mathbf{7}(3.2138, 0.8271)$, $\mathbf{15/2}(3.2503, 0.8327)$, $\mathbf{8}(3.2840, 0.8288)$.	0.368

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(2)}$	$ \mathcal{H}_{\text{CF}}^{(2)} _{\text{av}}$
8	$(0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ $\mathbf{1}(2.4495, 0.8164), \mathbf{3}/\mathbf{2}(2.0000, 1.0000), \mathbf{2}(2.3905, 0.8920), \mathbf{5}/\mathbf{2}(2.4495, 0.8164)$ $\mathbf{3}(2.6332, 0.8354), \mathbf{7}/\mathbf{2}(2.7373, 0.8627), \mathbf{4}(2.8470, 0.8439), \mathbf{9}/\mathbf{2}(2.9310, 0.8238)$ $\mathbf{5}(3.0058, 0.8299), \mathbf{11}/\mathbf{2}(3.0682, 0.8404), \mathbf{6}(3.1233, 0.8328), \mathbf{13}/\mathbf{2}(3.1709, 0.8241)$ $\mathbf{7}(3.2138, 0.8271), \mathbf{15}/\mathbf{2}(3.2503, 0.8327), \mathbf{8}(3.2840, 0.8288).$	0.368

No.	$\mathcal{H}_{\text{CF}}^{(4)}$	$ \mathcal{H}_{\text{CF}}^{(4)} _{\text{av}}$
1	$(1, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{2}(2.6725, 0.8553), \mathbf{5}/\mathbf{2}(2.3148, 1.9259), \mathbf{3}(2.7717, 0.8525), \mathbf{7}/\mathbf{2}(2.5074, 0.9113)$ $\mathbf{4}(2.6154, 0.9540), \mathbf{9}/\mathbf{2}(2.3651, 0.9221), \mathbf{5}(2.3531, 0.9273), \mathbf{11}/\mathbf{2}(2.5544, 0.9415)$ $\mathbf{6}(2.6944, 0.9183), \mathbf{13}/\mathbf{2}(2.7883, 0.9103), \mathbf{7}(2.8478, 0.9144), \mathbf{15}/\mathbf{2}(2.8812, 0.8952)$ $\mathbf{8}(2.8957, 0.8832).$	axial 0.287
2	$(0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0)$ $\mathbf{2}(2.7598, 0.8560), \mathbf{5}/\mathbf{2}(2.3780, 0.9061), \mathbf{3}(3.0011, 0.8326), \mathbf{7}/\mathbf{2}(2.6228, 0.8977)$ $\mathbf{4}(2.6514, 0.9396), \mathbf{9}/\mathbf{2}(2.4258, 0.9164), \mathbf{5}(2.3531, 0.9273), \mathbf{11}/\mathbf{2}(2.5222, 0.9457)$ $\mathbf{6}(2.6598, 0.9205), \mathbf{13}/\mathbf{2}(2.7423, 0.9081), \mathbf{7}(2.8060, 0.9086), \mathbf{15}/\mathbf{2}(2.8416, 0.8892)$ $\mathbf{8}(2.8684, 0.8757).$	0.283
3	$(\frac{1}{\sqrt{2}}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0)$ $\mathbf{2}(2.8651, 0.8426), \mathbf{5}/\mathbf{2}(2.3544, 0.9149), \mathbf{3}(2.7479, 0.8429), \mathbf{7}/\mathbf{2}(2.4981, 0.9215)$ $\mathbf{4}(2.5875, 0.9522), \mathbf{9}/\mathbf{2}(2.3679, 0.9193), \mathbf{5}(2.3531, 0.9233), \mathbf{11}/\mathbf{2}(2.5472, 0.9343)$ $\mathbf{6}(2.6836, 0.9108), \mathbf{13}/\mathbf{2}(2.7782, 0.9014), \mathbf{7}(2.8350, 0.9005), \mathbf{15}/\mathbf{2}(2.8848, 0.8844)$ $\mathbf{8}(2.9167, 0.8733).$	0.283
4	$(\frac{1}{2}\sqrt{\frac{7}{3}}, 0, 0, 0, 0, 0, \frac{1}{2}\sqrt{\frac{5}{6}}, \frac{1}{2}\sqrt{\frac{5}{6}})$ $\mathbf{2}(2.0412, 0.9801), \mathbf{5}/\mathbf{2}(2.1213, 0.9428), \mathbf{3}(2.9312, 0.8374), \mathbf{7}/\mathbf{2}(2.7849, 0.7832)$ $\mathbf{4}(2.7657, 0.8874), \mathbf{9}/\mathbf{2}(2.4467, 0.9458), \mathbf{5}(2.3531, 0.9681), \mathbf{11}/\mathbf{2}(2.3830, 0.9779)$ $\mathbf{6}(2.5333, 0.9746), \mathbf{13}/\mathbf{2}(2.4549, 0.9620), \mathbf{7}(2.4516, 0.9423), \mathbf{15}/\mathbf{2}(2.4636, 0.9204)$ $\mathbf{8}(2.5221, 0.8980).$	cubic 0.280
5	$(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, 0, 0, 0, 0, 0)$ $\mathbf{2}(2.9268, 0.8318), \mathbf{5}/\mathbf{2}(2.3824, 0.9039), \mathbf{3}(2.7336, 0.8588), \mathbf{7}/\mathbf{2}(2.5006, 0.9291)$ $\mathbf{4}(2.5650, 0.9513), \mathbf{9}/\mathbf{2}(2.3708, 0.9164), \mathbf{5}(2.3531, 0.9194), \mathbf{11}/\mathbf{2}(2.5388, 0.9280)$ $\mathbf{6}(2.6728, 0.9032), \mathbf{13}/\mathbf{2}(2.7670, 0.8935), \mathbf{7}(2.8339, 0.8947), \mathbf{15}/\mathbf{2}(2.8836, 0.8760)$ $\mathbf{8}(2.9340, 0.8646).$	0.280
6	$(0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2})$ $\mathbf{2}(2.5981, 0.9298), \mathbf{5}/\mathbf{2}(2.2935, 0.9303), \mathbf{3}(3.0249, 0.8072), \mathbf{7}/\mathbf{2}(2.6975, 0.8477)$ $\mathbf{4}(2.7594, 0.9180), \mathbf{9}/\mathbf{2}(2.4808, 0.9354), \mathbf{5}(2.3531, 0.9522), \mathbf{11}/\mathbf{2}(2.4692, 0.9623)$ $\mathbf{6}(2.6155, 0.9540), \mathbf{13}/\mathbf{2}(2.6412, 0.9418), \mathbf{7}(2.6944, 0.9191), \mathbf{15}/\mathbf{2}(2.7144, 0.8976)$ $\mathbf{8}(2.7472, 0.8881).$	0.279

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(4)}$	$ \mathcal{H}_{\text{CF}}^{(4)} _{\text{av}}$
7	$(\frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2})$ $\mathbf{2}(2.2358, 0.9760), \mathbf{5/2}(2.1715, 0.9421), \mathbf{3}(2.9558, 0.8350), \mathbf{7/2}(2.7781, 0.7815)$ $\mathbf{4}(2.7918, 0.8865), \mathbf{9/2}(2.4656, 0.9449), \mathbf{5}(2.3531, 0.9671), \mathbf{11/2}(2.3954, 0.9758)$ $\mathbf{6}(2.5419, 0.9735), \mathbf{13/2}(2.4774, 0.9609), \mathbf{7}(2.4725, 0.9365), \mathbf{15/2}(2.4924, 0.9192)$ $\mathbf{8}(2.5382, 0.8968).$	0.279
8	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}e^{i\pi/2}, \frac{1}{3}e^{-i\pi/2}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ $\mathbf{2}(2.8047, 0.8560), \mathbf{5/2}(2.3405, 0.9193), \mathbf{3}(3.1297, 0.8159), \mathbf{7/2}(2.7577, 0.8095)$ $\mathbf{4}(2.7846, 0.8937), \mathbf{9/2}(2.4799, 0.9335), \mathbf{5}(2.3531, 0.9273), \mathbf{11/2}(2.4266, 0.9675)$ $\mathbf{6}(2.6220, 0.9778), \mathbf{13/2}(2.4762, 0.9463), \mathbf{7}(2.6212, 0.9249), \mathbf{15/2}(2.6292, 0.9012)$ $\mathbf{8}(2.6780, 0.8819).$	0.277
9	$(\frac{1}{\sqrt{2}}, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0)$ $\mathbf{2}(2.3385, 0.9633), \mathbf{5/2}(2.3544, 0.9149), \mathbf{3}(3.0717, 0.8255), \mathbf{7/2}(2.7959, 0.7637)$ $\mathbf{4}(2.7450, 0.8703), \mathbf{9/2}(2.4590, 0.9345), \mathbf{5}(2.3531, 0.9631), \mathbf{11/2}(2.3996, 0.9738)$ $\mathbf{6}(2.5441, 0.9681), \mathbf{13/2}(2.5279, 0.9530), \mathbf{7}(2.5504, 0.9284), \mathbf{15/2}(2.5752, 0.9084)$ $\mathbf{8}(2.6248, 0.8844).$	0.277
10	$(\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ $\mathbf{2}(2.7289, 0.9103), \mathbf{5/2}(2.3736, 0.9075), \mathbf{3}(3.1662, 0.8239), \mathbf{7/2}(2.7891, 0.7789)$ $\mathbf{4}(2.7558, 0.8766), \mathbf{9/2}(2.4751, 0.9297), \mathbf{5}(2.3531, 0.9592), \mathbf{11/2}(2.4058, 0.9706)$ $\mathbf{6}(2.6263, 0.9908), \mathbf{13/2}(2.5357, 0.9474), \mathbf{7}(2.5794, 0.9249), \mathbf{15/2}(2.5800, 0.9012)$ $\mathbf{8}(2.6211, 0.8782).$	0.276
11	$(0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0)$ $\mathbf{2}(2.7530, 0.8989), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(3.0733, 0.7874), \mathbf{7/2}(2.7314, 0.8663)$ $\mathbf{4}(2.6550, 0.8991), \mathbf{9/2}(2.4286, 0.8975), \mathbf{5}(2.3531, 0.9423), \mathbf{11/2}(2.4630, 0.9530)$ $\mathbf{6}(2.5971, 0.9389), \mathbf{13/2}(2.6558, 0.9126), \mathbf{7}(3.0047, 0.9028), \mathbf{15/2}(2.7684, 0.8820)$ $\mathbf{8}(2.8190, 0.8597).$	0.276
12	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, -\frac{1}{3}, -\frac{1}{3})$ $\mathbf{2}(2.5438, 0.9284), \mathbf{5/2}(2.4110, 0.8870), \mathbf{3}(3.2860, 0.8207), \mathbf{7/2}(2.8035, 0.7705)$ $\mathbf{4}(2.7873, 0.8667), \mathbf{9/2}(2.4789, 0.9212), \mathbf{5}(2.3531, 0.9562), \mathbf{11/2}(2.4017, 0.9706)$ $\mathbf{6}(2.6122, 0.9876), \mathbf{13/2}(2.5312, 0.9418), \mathbf{7}(2.5980, 0.9179), \mathbf{15/2}(2.5716, 0.8940)$ $\mathbf{8}(2.6062, 0.8708).$	0.274
13	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, -\frac{1}{3}e^{i\pi/2}, -\frac{1}{3}e^{-i\pi/2}, \frac{1}{3}, \frac{1}{3})$ $\mathbf{2}(2.9228, 0.8358), \mathbf{5/2}(2.4272, 0.8730), \mathbf{3}(3.2289, 0.8128), \mathbf{7/2}(2.7789, 0.8171)$ $\mathbf{4}(2.7612, 0.8838), \mathbf{9/2}(2.4761, 0.9136), \mathbf{5}(2.3531, 0.9482), \mathbf{11/2}(2.4224, 0.9644)$ $\mathbf{6}(2.6317, 0.9767), \mathbf{13/2}(2.5705, 0.9306), \mathbf{7}(2.6352, 0.9063), \mathbf{15/2}(2.6436, 0.8832)$ $\mathbf{8}(2.6928, 0.8671).$	0.273

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(4)}$	$ \mathcal{H}_{\text{CF}}^{(4)} _{\text{av}}$
14	$(0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2})$ $\mathbf{2}(2.4961, 0.8996), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(3.3225, 0.7921), \mathbf{7/2}(2.8044, 0.7934)$ $\mathbf{4}(2.7774, 0.8676), \mathbf{9/2}(2.4837, 0.8918), \mathbf{5}(2.3531, 0.9423), \mathbf{11/2}(2.4287, 0.9675)$ $\mathbf{6}(2.5744, 0.9465), \mathbf{13/2}(2.5862, 0.9182), \mathbf{7}(2.6538, 0.9016), \mathbf{15/2}(2.6640, 0.8880)$ $\mathbf{8}(2.7064, 0.8659).$	0.273
15	$(\frac{3}{2\sqrt{11}}, 0, 0, 0, 0, 0, 0, \frac{1}{2}\sqrt{\frac{35}{22}}, \frac{1}{2}\sqrt{\frac{35}{22}})$ $\mathbf{2}(2.8208, 0.9023), \mathbf{5/2}(2.3398, 0.9193), \mathbf{3}(2.8701, 0.9048), \mathbf{7/2}(2.6109, 0.8935)$ $\mathbf{4}(2.7198, 0.9324), \mathbf{9/2}(2.4542, 0.9307), \mathbf{5}(2.3531, 0.9432), \mathbf{11/2}(2.4463, 0.9447)$ $\mathbf{6}(2.5798, 0.9346), \mathbf{13/2}(2.5829, 0.9216), \mathbf{7}(2.6480, 0.8993), \mathbf{15/2}(2.6676, 0.8772)$ $\mathbf{8}(2.7039, 0.8572).$	plane square 0.271
16	$(0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0, 0, 0)$ $\mathbf{2}(2.3908, 0.8922), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(3.3654, 0.7945), \mathbf{7/2}(2.8222, 0.7526)$ $\mathbf{4}(2.7999, 0.8424), \mathbf{9/2}(2.4922, 0.8880), \mathbf{5}(2.3531, 0.9393), \mathbf{11/2}(2.4141, 0.9706)$ $\mathbf{6}(2.5657, 0.9454), \mathbf{13/2}(2.5537, 0.9171), \mathbf{7}(2.6224, 0.9016), \mathbf{15/2}(2.6016, 0.8856)$ $\mathbf{8}(2.6260, 0.8634).$	0.269
17	$(0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0)$ $\mathbf{2}(2.2922, 0.8942), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(3.3821, 0.8048), \mathbf{7/2}(2.8281, 0.7238)$ $\mathbf{4}(2.8152, 0.8226), \mathbf{9/2}(2.4941, 0.8880), \mathbf{5}(2.3531, 0.9403), \mathbf{11/2}(2.4058, 0.9717)$ $\mathbf{6}(2.5365, 0.9465), \mathbf{13/2}(2.5312, 0.9171), \mathbf{7}(2.9710, 0.8888), \mathbf{15/2}(2.5680, 0.8700)$ $\mathbf{8}(2.5901, 0.8535).$	0.268
18	$(0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0)$ $\mathbf{2}(2.2358, 0.8942), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(3.3844, 0.8056), \mathbf{7/2}(2.8281, 0.7068)$ $\mathbf{4}(2.8161, 0.8118), \mathbf{9/2}(2.4950, 0.8880), \mathbf{5}(2.3531, 0.9403), \mathbf{11/2}(2.4037, 0.9717)$ $\mathbf{6}(2.5343, 0.9465), \mathbf{13/2}(2.5279, 0.9171), \mathbf{7}(2.6131, 0.8888), \mathbf{15/2}(2.5620, 0.8616)$ $\mathbf{8}(2.5827, 0.8535).$	0.266
19	$(0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, 0)$ $\mathbf{2}(3.0475, 0.7788), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(2.7114, 0.8699), \mathbf{7/2}(2.5312, 0.9487)$ $\mathbf{4}(2.5101, 0.9270), \mathbf{9/2}(2.3802, 0.8927), \mathbf{5}(2.3531, 0.9015), \mathbf{11/2}(2.5014, 0.9062)$ $\mathbf{6}(2.6230, 0.8794), \mathbf{13/2}(2.7176, 0.8565), \mathbf{7}(2.8071, 0.8447), \mathbf{15/2}(2.8956, 0.8268)$ $\mathbf{8}(2.9822, 0.8127).$	0.266
20	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ $\mathbf{2}(2.6035, 0.9090), \mathbf{5/2}(2.4478, 0.8333), \mathbf{3}(2.7645, 0.8977), \mathbf{7/2}(2.4802, 0.9588)$ $\mathbf{4}(2.4714, 0.9351), \mathbf{9/2}(2.3613, 0.8984), \mathbf{5}(2.3531, 0.8975), \mathbf{11/2}(2.4692, 0.8917)$ $\mathbf{6}(2.6890, 0.8956), \mathbf{13/2}(2.6513, 0.8475), \mathbf{7}(2.7305, 0.8389), \mathbf{15/2}(2.7840, 0.8316)$ $\mathbf{8}(2.8474, 0.8226).$	0.265

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(4)}$	$ \mathcal{H}_{\text{CF}}^{(4)} _{\text{av}}$
21	$(0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ $\mathbf{2}(3.0542, 0.7748), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(2.8169, 0.8977), \mathbf{7/2}(2.5227, 0.9504)$ $\mathbf{4}(2.5326, 0.9261), \mathbf{9/2}(2.3765, 0.8927), \mathbf{5}(2.3531, 0.9005), \mathbf{11/2}(2.4827, 0.9021)$ $\mathbf{6}(2.6014, 0.8783), \mathbf{13/2}(2.6794, 0.8576), \mathbf{7}(2.7595, 0.8377), \mathbf{15/2}(2.8284, 0.8196)$ $\mathbf{8}(2.8994, 0.8151).$	0.264
22	$(\frac{1}{\sqrt{3}}, 0, 0, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, 0, 0, 0, 0)$ $\mathbf{2}(3.0623, 0.7929), \mathbf{5/2}(2.4228, 0.8774), \mathbf{3}(2.7606, 0.9334), \mathbf{7/2}(2.4149, 0.9554)$ $\mathbf{4}(2.5056, 0.9477), \mathbf{9/2}(2.3736, 0.9126), \mathbf{5}(2.3531, 0.9064), \mathbf{11/2}(2.4723, 0.8969)$ $\mathbf{6}(2.5895, 0.8772), \mathbf{13/2}(2.6424, 0.8632), \mathbf{7}(2.7200, 0.8505), \mathbf{15/2}(2.7708, 0.8376)$ $\mathbf{8}(2.8350, 0.8287).$	0.263
23	$(\frac{1}{3}, \frac{1}{3}e^{i\pi/4}, \frac{1}{3}e^{-i\pi/4}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ $\mathbf{2}(2.7779, 0.8573), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(2.5748, 0.8969), \mathbf{7/2}(2.4353, 0.9682)$ $\mathbf{4}(2.4300, 0.9342), \mathbf{9/2}(2.3395, 0.8937), \mathbf{5}(2.3531, 0.8895), \mathbf{11/2}(2.4910, 0.8771)$ $\mathbf{6}(2.6717, 0.8805), \mathbf{13/2}(2.6929, 0.8295), \mathbf{7}(2.7804, 0.8215), \mathbf{15/2}(2.7660, 0.8136)$ $\mathbf{8}(2.9464, 0.7966).$	0.257
24	$(\frac{1}{\sqrt{2}}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0)$ $\mathbf{2}(3.1334, 0.7245), \mathbf{5/2}(2.4412, 0.8532), \mathbf{3}(2.5336, 0.9342), \mathbf{7/2}(2.3326, 0.9750)$ $\mathbf{4}(2.3625, 0.9486), \mathbf{9/2}(2.3138, 0.9050), \mathbf{5}(2.3531, 0.8835), \mathbf{11/2}(2.4806, 0.8626)$ $\mathbf{6}(2.5938, 0.8372), \mathbf{13/2}(2.6570, 0.8284), \mathbf{7}(2.7363, 0.8203), \mathbf{15/2}(2.7972, 0.8160)$ $\mathbf{8}(2.8660, 0.8139).$	0.257
25	$(0, 0, 0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ $\mathbf{2}(3.1623, 0.6326), \mathbf{5/2}(2.4495, 0.8164), \mathbf{3}(2.5225, 0.9183), \mathbf{7/2}(2.3351, 0.9826)$ $\mathbf{4}(2.3796, 0.9396), \mathbf{9/2}(2.2901, 0.8946), \mathbf{5}(2.3531, 0.8527), \mathbf{11/2}(2.4838, 0.8168)$ $\mathbf{6}(2.5960, 0.8004), \mathbf{13/2}(2.6614, 0.7195), \mathbf{7}(2.7386, 0.7971), \mathbf{15/2}(2.8020, 0.8028)$ $\mathbf{8}(2.8684, 0.7991).$	0.251

No.	$\mathcal{H}_{\text{CF}}^{(6)}$	$ \mathcal{H}_{\text{CF}}^{(6)} _{\text{av}}$
1	$(1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(3.0459, 0.7956), \mathbf{7/2}(2.4373, 0.8709), \mathbf{4}(2.8318, 0.8091), \mathbf{9/2}(2.5848, 0.9361)$ $\mathbf{5}(2.6296, 0.9112), \mathbf{11/2}(2.8952, 0.8793), \mathbf{6}(2.9640, 0.8840), \mathbf{13/2}(2.8965, 0.8418)$ $\mathbf{7}(2.8655, 0.8951), \mathbf{15/2}(2.9306, 0.9201), \mathbf{8}(2.9004, 0.9009).$	axial 0.239
2	$(\frac{1}{\sqrt{7}}, 0, 0, \frac{1}{\sqrt{7}}, \frac{1}{\sqrt{7}}, 0, 0, \frac{1}{\sqrt{7}}, \frac{1}{\sqrt{7}}, 0, 0, \frac{1}{\sqrt{7}}, \frac{1}{\sqrt{7}})$ $\mathbf{3}(3.2615, 0.7832), \mathbf{7/2}(2.6117, 0.8597), \mathbf{4}(3.0330, 0.8231), \mathbf{9/2}(2.5905, 0.9315)$ $\mathbf{5}(2.6260, 0.9052), \mathbf{11/2}(3.0763, 0.8243), \mathbf{6}(2.9250, 0.8723), \mathbf{13/2}(2.8627, 0.8526)$ $\mathbf{7}(2.8431, 0.8979), \mathbf{15/2}(2.9032, 0.9173), \mathbf{8}(2.8751, 0.8994).$	0.236

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(6)}$	$ \mathcal{H}_{\text{CF}}^{(6)} _{\text{av}}$
3	$(\frac{1}{\sqrt{2}}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(3.2672, 0.7832), \mathbf{7/2}(2.6229, 0.8587), \mathbf{4}(2.8286, 0.8275), \mathbf{9/2}(2.5893, 0.9304)$ $\mathbf{5}(2.6201, 0.9040), \mathbf{11/2}(2.8602, 0.8706), \mathbf{6}(2.9289, 0.8723), \mathbf{13/2}(2.8708, 0.8526)$ $\mathbf{7}(2.8375, 0.8979), \mathbf{15/2}(2.8974, 0.9173), \mathbf{8}(2.8706, 0.8979).$	0.235
4	$(0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(3.1995, 0.7870), \mathbf{7/2}(2.6352, 0.8689), \mathbf{4}(3.0806, 0.8448), \mathbf{9/2}(2.6520, 0.9372)$ $\mathbf{5}(2.8329, 0.9100), \mathbf{11/2}(2.9164, 0.8543), \mathbf{6}(2.9906, 0.8645), \mathbf{13/2}(2.9410, 0.8418)$ $\mathbf{7}(2.9213, 0.8881), \mathbf{15/2}(2.9465, 0.9115), \mathbf{8}(2.9286, 0.9053).$	0.234
5	$(0, 0, 0, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, 0, 0, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, 0, 0, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}})$ $\mathbf{3}(2.9047, 0.8738), \mathbf{7/2}(2.4934, 0.9137), \mathbf{4}(3.1595, 0.7810), \mathbf{9/2}(2.6019, 0.9224)$ $\mathbf{5}(2.8544, 0.9136), \mathbf{11/2}(2.9889, 0.8431), \mathbf{6}(3.3176, 0.8437), \mathbf{13/2}(3.1150, 0.8607)$ $\mathbf{7}(3.2495, 0.8532), \mathbf{15/2}(3.0618, 0.8999), \mathbf{8}(2.9613, 0.9053).$	0.233
6	$(\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}})$ $\mathbf{3}(2.4459, 0.9740), \mathbf{7/2}(2.5679, 0.8362), \mathbf{4}(2.6869, 0.9616), \mathbf{9/2}(2.3134, 0.9657)$ $\mathbf{5}(2.7384, 0.8287), \mathbf{11/2}(2.9864, 0.8481), \mathbf{6}(3.4606, 0.8762), \mathbf{13/2}(3.1663, 0.8715)$ $\mathbf{7}(3.0372, 0.8323), \mathbf{15/2}(2.8397, 0.8019), \mathbf{8}(2.9360, 0.8310).$	0.231
7	$(0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(3.0097, 0.8376), \mathbf{7/2}(2.7555, 0.8485), \mathbf{4}(3.1011, 0.8826), \mathbf{9/2}(2.5825, 0.9087)$ $\mathbf{5}(2.8628, 0.8598), \mathbf{11/2}(3.0251, 0.8268), \mathbf{6}(3.2591, 0.8411), \mathbf{13/2}(3.1231, 0.8216)$ $\mathbf{7}(3.1350, 0.8393), \mathbf{15/2}(2.9840, 0.8826), \mathbf{8}(3.0148, 0.8726).$	0.231
8	$(0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2})$ $\mathbf{3}(2.8456, 0.8957), \mathbf{7/2}(2.5587, 0.9413), \mathbf{4}(3.3175, 0.7258), \mathbf{9/2}(2.6714, 0.8916)$ $\mathbf{5}(2.8975, 0.8849), \mathbf{11/2}(3.0576, 0.7856), \mathbf{6}(3.4697, 0.8307), \mathbf{13/2}(3.2391, 0.8364)$ $\mathbf{7}(3.3109, 0.8295), \mathbf{15/2}(3.1238, 0.8610), \mathbf{8}(2.9509, 0.8771).$	0.230
9	$(\frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2})$ $\mathbf{3}(2.6462, 0.9234), \mathbf{7/2}(2.4200, 0.9637), \mathbf{4}(3.3402, 0.7377), \mathbf{9/2}(2.6156, 0.9133)$ $\mathbf{5}(2.9692, 0.9148), \mathbf{11/2}(3.0563, 0.8668), \mathbf{6}(3.3670, 0.8112), \mathbf{13/2}(3.2432, 0.8486)$ $\mathbf{7}(3.4366, 0.8462), \mathbf{15/2}(3.1354, 0.8624), \mathbf{8}(3.0193, 0.8949).$	0.229
10	$(\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}})$ $\mathbf{3}(2.7273, 0.8948), \mathbf{7/2}(2.5597, 0.9403), \mathbf{4}(3.4191, 0.8015), \mathbf{9/2}(2.6042, 0.8973)$ $\mathbf{5}(3.0625, 0.9088), \mathbf{11/2}(3.0713, 0.8755), \mathbf{6}(3.6491, 0.8879), \mathbf{13/2}(3.2067, 0.8337)$ $\mathbf{7}(3.3444, 0.8379), \mathbf{15/2}(3.1065, 0.8639), \mathbf{8}(3.0446, 0.8920).$	0.228
11	$(\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}})$ $\mathbf{3}(2.8303, 0.9206), \mathbf{7/2}(2.7494, 0.8536), \mathbf{4}(2.9724, 0.8426), \mathbf{9/2}(2.4936, 0.8939)$ $\mathbf{5}(2.8999, 0.8490), \mathbf{11/2}(3.1100, 0.8143), \mathbf{6}(3.6387, 0.8866), \mathbf{13/2}(3.2432, 0.8364)$ $\mathbf{7}(3.2132, 0.8127), \mathbf{15/2}(2.9984, 0.8480), \mathbf{8}(2.8840, 0.8697).$	0.227

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(6)}$	$ \mathcal{H}_{\text{CF}}^{(6)} _{\text{av}}$
12	$(\frac{1}{\sqrt{3}}, 0, 0, 0, 0, 0, 0, 0, 0, 0, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$ $\mathbf{3}(3.0555, 0.8814), \mathbf{7/2}(2.3874, 0.9484), \mathbf{4}(3.3553, 0.7853), \mathbf{9/2}(2.6258, 0.9247)$ $\mathbf{5}(2.9489, 0.9339), \mathbf{11/2}(2.9889, 0.9005), \mathbf{6}(3.2487, 0.8515), \mathbf{13/2}(3.1285, 0.8688)$ $\mathbf{7}(3.3305, 0.8686), \mathbf{15/2}(3.0763, 0.8999), \mathbf{8}(3.0312, 0.9143).$	0.226
13	$(-\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}})$ $\mathbf{3}(3.3226, 0.7937), \mathbf{7/2}(2.7504, 0.8505), \mathbf{4}(3.1325, 0.8653), \mathbf{9/2}(2.5643, 0.8871)$ $\mathbf{5}(2.8568, 0.8909), \mathbf{11/2}(2.9264, 0.8743), \mathbf{6}(3.5243, 0.9048), \mathbf{13/2}(3.0071, 0.8567)$ $\mathbf{7}(2.9870, 0.8770), \mathbf{15/2}(2.9090, 0.8855), \mathbf{8}(2.8974, 0.8786).$	0.225
14	$(\frac{1}{\sqrt{5}}, 0, 0, 0, 0, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, 0, 0, 0, 0, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}})$ $\mathbf{3}(2.7197, 0.9406), \mathbf{7/2}(2.6790, 0.8923), \mathbf{4}(3.3737, 0.7615), \mathbf{9/2}(2.5563, 0.9087)$ $\mathbf{5}(2.6906, 0.8861), \mathbf{11/2}(3.0001, 0.8331), \mathbf{6}(3.2981, 0.8710), \mathbf{13/2}(3.2459, 0.8594)$ $\mathbf{7}(3.2537, 0.8518), \mathbf{15/2}(2.9998, 0.8596), \mathbf{8}(2.8186, 0.8786).$	0.224
15	$(\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}})$ $\mathbf{3}(3.1442, 0.8347), \mathbf{7/2}(2.6444, 0.9056), \mathbf{4}(3.3315, 0.8307), \mathbf{9/2}(2.6042, 0.9076)$ $\mathbf{5}(3.0314, 0.9124), \mathbf{11/2}(2.9789, 0.8843), \mathbf{6}(3.5178, 0.8892), \mathbf{13/2}(3.0907, 0.8513)$ $\mathbf{7}(3.2048, 0.8630), \mathbf{15/2}(3.0301, 0.8841), \mathbf{8}(3.0802, 0.8860).$	0.223
16	$(\frac{1}{2\sqrt{2}}, 0, 0, 0, 0, 0, 0, -\frac{\sqrt{7}}{4}, -\frac{\sqrt{7}}{4}, 0, 0, 0, 0)$ $\mathbf{3}(2.5852, 0.9492), \mathbf{7/2}(2.2160, 0.9851), \mathbf{4}(3.4321, 0.7420), \mathbf{9/2}(2.5437, 0.9201)$ $\mathbf{5}(2.6033, 0.8885), \mathbf{11/2}(2.8565, 0.8318), \mathbf{6}(3.2929, 0.8099), \mathbf{13/2}(3.2486, 0.8567)$ $\mathbf{7}(3.2565, 0.8588), \mathbf{15/2}(2.9782, 0.8596), \mathbf{8}(2.7889, 0.8816).$	cubic 0.223
17	$(\frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0)$ $\mathbf{3}(2.6023, 0.8910), \mathbf{7/2}(2.4271, 0.9668), \mathbf{4}(3.5879, 0.7669), \mathbf{9/2}(2.6498, 0.8882)$ $\mathbf{5}(3.0745, 0.9017), \mathbf{11/2}(3.0488, 0.8755), \mathbf{6}(3.3228, 0.8112), \mathbf{13/2}(3.1960, 0.8297)$ $\mathbf{7}(3.2565, 0.8113), \mathbf{15/2}(3.1065, 0.8437), \mathbf{8}(3.0208, 0.8771).$	0.223
18	$(\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}}, -\frac{1}{\sqrt{13}})$ $\mathbf{3}(3.1490, 0.8347), \mathbf{7/2}(2.7055, 0.8811), \mathbf{4}(3.2731, 0.8545), \mathbf{9/2}(2.5814, 0.8825)$ $\mathbf{5}(3.0434, 0.8993), \mathbf{11/2}(3.0113, 0.8868), \mathbf{6}(3.6894, 0.8658), \mathbf{13/2}(3.0934, 0.8472)$ $\mathbf{7}(3.1503, 0.8588), \mathbf{15/2}(2.9984, 0.8798), \mathbf{8}(2.9613, 0.8830).$	0.222
19	$(0, 0, 0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2})$ $\mathbf{3}(3.1509, 0.8013), \mathbf{7/2}(2.7463, 0.8556), \mathbf{4}(3.0795, 0.8286), \mathbf{9/2}(2.5403, 0.8985)$ $\mathbf{5}(2.9453, 0.8993), \mathbf{11/2}(2.9889, 0.8381), \mathbf{6}(3.1967, 0.8060), \mathbf{13/2}(3.0732, 0.8256)$ $\mathbf{7}(3.1224, 0.8546), \mathbf{15/2}(3.0099, 0.8798), \mathbf{8}(2.9717, 0.8935).$	0.222
20	$(0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(2.6809, 0.7777), \mathbf{7/2}(2.6713, 0.9003), \mathbf{4}(3.3986, 0.8664), \mathbf{9/2}(2.5807, 0.8817)$ $\mathbf{5}(3.1264, 0.8965), \mathbf{11/2}(3.0680, 0.8894), \mathbf{6}(3.1789, 0.8280), \mathbf{13/2}(3.1162, 0.7826)$ $\mathbf{7}(3.1029, 0.7905), \mathbf{15/2}(3.0760, 0.8365), \mathbf{8}(3.0453, 0.8659).$	0.221

Appendix – cont.

No.	$\mathcal{H}_{\text{CF}}^{(6)}$	$ \mathcal{H}_{\text{CF}}^{(6)} _{\text{av}}$
21	$(0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0)$ $\mathbf{3}(2.3934, 0.9120), \mathbf{7/2}(2.8284, 0.7071), \mathbf{4}(2.5041, 0.8156), \mathbf{9/2}(2.3351, 0.8939)$ $\mathbf{5}(2.7121, 0.7857), \mathbf{11/2}(3.1849, 0.7981), \mathbf{6}(3.6777, 0.8190), \mathbf{13/2}(3.3025, 0.8121)$ $\mathbf{7}(3.0819, 0.7890), \mathbf{15/2}(2.8888, 0.7903), \mathbf{8}(2.7651, 0.8280).$	0.221
22	$(0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0)$ $\mathbf{3}(2.6691, 0.8442), \mathbf{7/2}(2.5322, 0.9484), \mathbf{4}(3.5241, 0.8134), \mathbf{9/2}(2.6315, 0.8768)$ $\mathbf{5}(3.1247, 0.9040), \mathbf{11/2}(3.0538, 0.8880), \mathbf{6}(3.3670, 0.8125), \mathbf{13/2}(3.1609, 0.8108)$ $\mathbf{7}(3.3039, 0.8267), \mathbf{15/2}(3.1123, 0.8524), \mathbf{8}(3.0357, 0.8667).$	0.221
23	$(0, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, 0, 0, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, 0, 0, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, 0, 0)$ $\mathbf{3}(3.0431, 0.8223), \mathbf{7/2}(2.8187, 0.7638), \mathbf{4}(2.8556, 0.8751), \mathbf{9/2}(3.0500, 0.8586)$ $\mathbf{5}(3.0159, 0.8454), \mathbf{11/2}(3.1400, 0.8281), \mathbf{6}(3.5555, 0.8099), \mathbf{13/2}(3.2270, 0.7987)$ $\mathbf{7}(3.1713, 0.8113), \mathbf{15/2}(2.9666, 0.8206), \mathbf{8}(2.9375, 0.8399).$	0.220
24	$(0, 0, 0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0, 0, 0)$ $\mathbf{3}(2.7634, 0.9044), \mathbf{7/2}(2.3356, 0.9828), \mathbf{4}(3.6181, 0.7298), \mathbf{9/2}(2.6630, 0.8737)$ $\mathbf{5}(2.7589, 0.8352), \mathbf{11/2}(3.0346, 0.7842), \mathbf{6}(3.5205, 0.7887), \mathbf{13/2}(3.2465, 0.8359)$ $\mathbf{7}(3.3824, 0.8334), \mathbf{15/2}(3.0871, 0.8450), \mathbf{8}(2.9640, 0.8673).$	0.219
25	$(0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(3.4943, 0.7288), \mathbf{7/2}(2.7932, 0.8097), \mathbf{4}(2.8761, 0.8405), \mathbf{9/2}(2.5996, 0.8802)$ $\mathbf{5}(2.6356, 0.8490), \mathbf{11/2}(2.7690, 0.8643), \mathbf{6}(2.8418, 0.8762), \mathbf{13/2}(2.8573, 0.8715)$ $\mathbf{7}(2.8585, 0.8923), \mathbf{15/2}(2.8484, 0.8999), \mathbf{8}(2.8246, 0.8830).$	0.219
26	$(\frac{1}{\sqrt{5}}, 0, 0, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, 0, 0, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, 0, 0, 0, 0)$ $\mathbf{3}(3.5153, 0.7612), \mathbf{7/2}(2.7902, 0.8036), \mathbf{4}(2.9930, 0.8340), \mathbf{9/2}(2.5985, 0.8928)$ $\mathbf{5}(2.6571, 0.8801), \mathbf{11/2}(2.7753, 0.8843), \mathbf{6}(2.9523, 0.8905), \mathbf{13/2}(2.8492, 0.8796)$ $\mathbf{7}(2.7998, 0.8839), \mathbf{15/2}(2.7503, 0.8870), \mathbf{8}(2.7041, 0.8682).$	0.216
27	$(\frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{1}{\sqrt{13}})$ $\mathbf{3}(3.5964, 0.7221), \mathbf{7/2}(2.8106, 0.7801), \mathbf{4}(2.9216, 0.8405), \mathbf{9/2}(2.6110, 0.8802)$ $\mathbf{5}(2.7647, 0.9040), \mathbf{11/2}(2.7428, 0.9205), \mathbf{6}(3.0797, 0.9490), \mathbf{13/2}(2.7858, 0.8931)$ $\mathbf{7}(2.8319, 0.8909), \mathbf{15/2}(2.7272, 0.8783), \mathbf{8}(2.8052, 0.8667).$	0.213
28	$(0, 0, 0, 0, 0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0)$ $\mathbf{3}(2.6462, 0.7555), \mathbf{7/2}(2.8289, 0.7067), \mathbf{4}(2.6836, 0.8945), \mathbf{9/2}(2.4491, 0.8893)$ $\mathbf{5}(3.1151, 0.8730), \mathbf{11/2}(3.0875, 0.7769), \mathbf{6}(3.3267, 0.7761), \mathbf{13/2}(3.1339, 0.7393)$ $\mathbf{7}(3.1964, 0.7848), \mathbf{15/2}(3.0027, 0.8235), \mathbf{8}(2.9881, 0.8578).$	0.212
29	$(\frac{1}{\sqrt{2}}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0, 0, 0, 0)$ $\mathbf{3}(3.6622, 0.6773), \mathbf{7/2}(2.8157, 0.7598), \mathbf{4}(2.8102, 0.8264), \mathbf{9/2}(2.6680, 0.9099)$ $\mathbf{5}(2.6464, 0.9136), \mathbf{11/2}(2.6129, 0.9318), \mathbf{6}(2.6533, 0.9256), \mathbf{13/2}(2.6334, 0.9052)$ $\mathbf{7}(2.6155, 0.8979), \mathbf{15/2}(2.5960, 0.8826), \mathbf{8}(2.5837, 0.8652).$	0.207
30	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ $\mathbf{3}(3.7414, 0.5342), \mathbf{7/2}(2.8289, 0.7067), \mathbf{4}(2.8978, 0.8091), \mathbf{9/2}(2.6452, 0.8757)$ $\mathbf{5}(2.6918, 0.9220), \mathbf{11/2}(2.5667, 0.9555), \mathbf{6}(2.6000, 0.9802), \mathbf{13/2}(2.5241, 0.9147)$ $\mathbf{7}(2.5499, 0.8895), \mathbf{15/2}(2.4979, 0.8639), \mathbf{8}(2.5183, 0.8399).$	0.195

Table 1: Extreme total splittings ΔE_{min} , ΔE_{max} , the averages $\overline{\Delta E}$ and the mean square deviations δ for $|J\rangle$ states exposed to the action of 8 representative $\mathcal{H}_{CF}^{(2)}$ s listed in Appendix, all values are given in σ . Please note, that in Tables 1 – 3 certain extreme values are given with the accuracy of $10^{-4}\sigma$, those found only in the representative $\mathcal{H}_{CF}^{(k)}$ sets with the accuracy of $10^{-2}\sigma$, but their averages and mean square deviations the accuracy of $10^{-3}\sigma$.

J	ΔE_{min} ΔE_{max}	No. of $\mathcal{H}_{CF}^{(2)}$ acc. to the list in Appendix	$\overline{\Delta E}$	δ
1	2.1213 2.4495	1 8,7	2.343	0.134
3/2	2.0000 2.0000	1 8,7	2.0000	0.0000
2	2.3905 2.3905	every	2.3905	0.0000
5/2	2.4056 2.4495	1 8,7	2.433	0.018
3	2.5983 2.6332	1 8,7	2.620	0.014
7/2	2.6184 2.7373	1 8,7	2.693	0.049
4	2.7356 2.8470	1 8,7	2.805	0.046
9/2	2.7540 2.9310	1 8,7	2.867	0.073
5	2.8308 3.0058	1 8,7	2.942	0.072
11/2	2.8443 3.0682	1 8,7	2.990	0.092
6	2.9008 3.1233	1 8,7	3.045	0.091
13/2	2.9124 3.1709	1 8,7	3.082	0.106
7	2.9549 3.2138	1 8,7	3.124	0.106
15/2	2.9641 3.2503	1 8,7	3.153	0.117
8	2.9973 3.2840	1 8,7	3.185	0.117

Table 2: Extreme total splittings ΔE_{min} , ΔE_{max} , the averages $\overline{\Delta E}$ and the mean square deviations δ for $|J\rangle$ states exposed to the action of 25 representative $\mathcal{H}_{CF}^{(4)}$ s listed in Appendix, all values are given in σ .

J	ΔE_{min} ΔE_{max}	No. of $\mathcal{H}_{CF}^{(4)}$ acc. to the list in Appendix	$\overline{\Delta E}$	δ
2	2.0412 3.1623	4 25	2.691	0.303
5/2	2.1213 2.4495	4 25	2.385	0.086
3	2.52 3.38	25 18	2.965	0.265
7/2	2.33 2.8284	24,25 18	2.640	0.167
4	2.36 2.82	24,25 18	2.654	0.144
9/2	2.29 2.50	25 18	2.423	0.062
5	2.3531 2.3531	every	2.3531	0.0000
11/2	2.38 2.55	4 1	2.456	0.050
6	2.53 2.69	4 1	2.608	0.0450
13/2	2.45 2.79	4 1	2.617	0.096
7	2.45 3.00	4 11	2.708	0.134
15/2	2.46 2.90	4 19	2.706	0.129
8	2.52 2.98	4 19	2.757	0.140

Table 3: Extreme total splittings ΔE_{min} , ΔE_{max} , the averages $\overline{\Delta E}$ and the mean square deviations δ for $|J\rangle$ states exposed to the action of 30 representative $\mathcal{H}_{CF}^{(6)}$ s listed in Appendix, all values are given in σ .

J	ΔE_{min} ΔE_{max}	No. of $\mathcal{H}_{CF}^{(4)}$ acc. to the list in Appendix	$\overline{\Delta E}$	δ
3	2.39 3.7417	21 30	3.004	0.363
7/2	2.22 2.8284	16 30,28,21	2.634	0.166
4	2.50 3.62	21 24	3.114	0.286
9/2	2.31 3.05	6,21 23	2.593	0.119
5	2.60 3.13	16 20,28,21	2.849	0.172
11/2	2.57 3.18	30 21	2.963	0.145
6	2.60 3.69	30 18	3.260	0.292
13/2	2.52 3.30	30 21	3.061	0.194
7	2.55 3.44	30 9	3.089	0.225
15/2	2.50 3.14	30 9	2.954	0.150
8	2.52 3.08	30 15,20	2.905	0.132

Table 4: Extreme total splittings of $|J\rangle$ states in virtual CF fields of Hamiltonians $\mathcal{H}_{\text{CF}}(q = 0) = B_{20}C_0^{(2)} + B_{40}C_0^{(4)} + B_{60}C_0^{(6)}$ yielding constant σ^2 in comparison with ΔE for the pure component 2^k -poles, all values are given in σ .

Quantum number	The total splitting ΔE in the 2^k -pole component			The range of the total splittings for the 2^k -pole superpositions	
J	$k = 2$	$k = 4$	$k = 6$	ΔE_{min}	ΔE_{max}
1	2.1213	—	—	2.1213	2.1213
3/2	2.0000	—	—	2.0000	2.0000
2	2.3906	2.6725	—	2.0412	2.7386
5/2	2.4056	2.3148	—	2.1213	2.4495
3	2.5984	2.7717	3.0461	2.0207	3.2404
7/2	2.6183	2.5074	2.4373	2.0000	2.8284
4	2.7351	2.6154	2.8317	2.0802	3.5025
9/2	2.7528	2.3651	2.5848	2.1820	3.1424
5	2.8307	2.3531	2.6302	2.2186	3.3534
11/2	2.8444	2.5544	2.8953	2.3075	3.3182
6	2.9007	2.6944	2.9641	2.3350	3.4603
13/2	2.9125	2.7883	2.8964	2.3710	3.4518
7	2.9551	2.8478	2.8656	2.4081	3.6067
15/2	2.9640	2.8812	2.9306	2.4293	3.6152
8	2.9975	2.8957	2.9004	2.4532	3.7450

Table 5: Extreme total splittings of $|J\rangle$ states in virtual CF fields of Hamiltonians $\mathcal{H}_{\text{CF}}(q = 1) = B_{21}C_1^{(2)} + B_{41}C_1^{(4)} + B_{61}C_1^{(6)}$ yielding constant σ^2 in comparison with ΔE for the pure component 2^k -poles, all values are given in σ .

Quantum number	The total splitting ΔE in the 2^k -pole component			The range of the total splittings for the 2^k -pole superpositions	
J	$k = 2$	$k = 4$	$k = 6$	ΔE_{min}	ΔE_{max}
1	2.4495	—	—	2.4495	2.4495
3/2	2.0000	—	—	2.0000	2.0000
2	2.3906	3.0475	—	2.2361	3.1623
5/2	2.4495	2.4495	—	2.4495	2.4495
3	2.6333	2.7114	3.4943	2.1605	3.7417
7/2	2.7373	2.5312	2.7932	2.0000	2.8284
4	2.8470	2.5101	2.8762	2.2354	3.4166
9/2	2.9310	2.3803	2.5996	2.2363	3.1623
5	3.0058	2.3531	2.6356	2.3534	3.4198
11/2	3.0682	2.5014	2.7690	2.4059	3.4636
6	3.1233	2.6230	2.8418	2.4835	3.6040
13/2	3.1709	2.7176	2.8573	2.5498	3.7381
7	3.2138	2.8071	2.8585	2.6229	3.8661
15/2	3.2504	2.8956	2.8484	2.6316	3.9880
8	3.2840	2.9822	2.8246	2.6695	4.1043

Table 6: Extreme total splittings of $|J\rangle$ states in virtual CF fields of Hamiltonians $\mathcal{H}_{\text{CF}}(q = 2) = B_{22}C_2^{(2)} + B_{42}C_4^{(4)} + B_{62}C_2^{(6)}$ yielding constant σ^2 in comparison with ΔE for the pure component 2^k -poles, all values are given in σ .

Quantum number	The total splitting ΔE in the 2^k -pole component			The range of the total splittings for the 2^k -pole superpositions	
J	$k = 2$	$k = 4$	$k = 6$	ΔE_{min}	ΔE_{max}
1	2.4495	—	—	2.4495	2.4495
3/2	2.0000	—	—	2.0000	2.0000
2	2.3906	2.3906	—	2.2362	3.1623
5/2	2.4495	2.4495	—	2.4495	2.4495
3	2.6333	3.3654	2.6809	2.1605	3.7417
7/2	2.7373	2.8222	2.6713	2.0000	2.8284
4	2.8470	2.7999	3.3986	2.2011	3.9518
9/2	2.9310	2.4922	2.5807	2.2361	3.1622
5	3.0058	2.3531	3.1263	2.3461	3.3533
11/2	3.0682	2.4141	3.0680	2.3249	3.4455
6	3.1233	2.5657	3.1788	2.4153	3.5953
13/2	3.1709	2.5537	3.1163	2.4521	3.6830
7	3.2138	2.6224	3.1028	2.5064	3.7913
15/2	3.2504	2.6016	3.0760	2.5015	3.8838
8	3.2840	2.6260	3.0454	2.5249	3.9745

Table 7: Extreme total splittings of $|J\rangle$ states in virtual CF fields of Hamiltonians $\mathcal{H}_{\text{CF}}(q = 3) = B_{43}C_3^{(4)} + B_{63}C_3^{(6)}$ yielding constant σ^2 in comparison with ΔE for the pure component 2^k -poles, all values are given in σ .

Quantum number	The total splitting ΔE in the 2^k -pole component		The range of the total splittings for the 2^k -pole superpositions	
J	$k = 4$	$k = 6$	ΔE_{min}	ΔE_{max}
2	2.2361	—	2.2361	2.2361
5/2	2.4495	—	2.4495	2.4495
3	3.3845	2.3934	2.1605	3.7417
7/2	2.8284	2.8284	2.8284	2.8284
4	2.8161	2.5041	2.4499	3.6586
9/2	2.4950	2.3351	2.2362	3.1622
5	2.3531	2.7121	2.3469	3.5904
11/2	2.4037	3.1850	2.3301	3.2130
6	2.5343	3.6777	2.4945	3.7054
13/2	2.5279	3.3025	2.4540	3.3101
7	2.6131	3.0819	2.4537	3.0833
15/2	2.5620	2.8888	2.4600	2.9024
8	2.5827	2.7651	2.5079	2.9333

Table 8: Extreme total splittings of $|J\rangle$ states in virtual CF fields of Hamiltonians $\mathcal{H}_{\text{CF}}(q = 4) = B_{44}C_4^{(4)} + B_{64}C_4^{(6)}$ yielding constant σ^2 in comparison with ΔE for the pure component 2^k -poles, all values are given in σ .

Quantum number	The total splitting ΔE in the 2^k -pole component		The range of the total splittings for the 2^k -pole superpositions	
J	$k = 4$	$k = 6$	ΔE_{min}	ΔE_{max}
2	3.1623	—	3.1623	3.1623
5/2	2.4495	—	2.4495	2.4495
3	2.5225	2.7634	2.1605	3.7417
7/2	2.3352	2.3355	2.0000	2.8284
4	2.3796	3.6181	2.1857	4.0763
9/2	2.2901	2.6629	2.2362	3.1623
5	2.3531	2.7589	2.3514	3.0557
11/2	2.4838	3.0346	2.4495	3.1697
6	2.5960	3.5206	2.3785	3.6621
13/2	2.6614	3.2465	2.4176	3.3513
7	2.7386	3.3824	2.4212	3.4501
15/2	2.8020	3.0871	2.3992	3.1517
8	2.8684	2.9640	2.3673	3.0352

Table 9: Total splitting ΔE (in σ) of $|J\rangle$ states in virtual CF fields of partial Hamiltonians $\mathcal{H}_{\text{CF}}^{(6)} = B_{65}C_5^{(6)}$ and $\mathcal{H}_{\text{CF}}^{(6)} = B_{66}C_6^{(6)}$ yielding the same σ^2 .

Quantum number J	ΔE	
	$q = 5$	$q = 6$
3	2.6458	3.7417
7/2	2.8284	2.8284
4	2.6835	2.8980
9/2	2.4495	2.6454
5	3.1152	2.6919
11/2	3.0872	2.5667
6	3.3269	2.6002
13/2	3.1339	2.5241
7	3.1964	2.5499
15/2	3.0030	2.4979
8	2.9881	2.5183

Table 10: Nominally extreme total splittings $\Delta\mathcal{E}$ (conserving σ^2) of states $|J\rangle$ expressed in σ .

Quantum number J	$\Delta\mathcal{E}_{min}$	$\Delta\mathcal{E}_{max}$
1	$3/\sqrt{2} = 2.1213$	$\sqrt{6} = 2.4495$
3/2	2.0000	2.0000
2	$5/\sqrt{6} = 2.0412$	$\sqrt{10} = 3.1623$
5/2	$3/\sqrt{2} = 2.1213$	$\sqrt{6} = 2.4495$
3	$7/\sqrt{12} = 2.0207$	$\sqrt{14} = 3.7417$
7/2	2.0000	$\sqrt{8} = 2.8284$
4	$9/\sqrt{20} = 2.0125$	$\sqrt{18} = 4.2426$
9/2	$5/\sqrt{6} = 2.0412$	$\sqrt{10} = 3.1623$
5	$11/\sqrt{30} = 2.0083$	$\sqrt{22} = 4.6904$
11/2	2.0000	$\sqrt{12} = 3.4641$
6	$13/\sqrt{42} = 2.0059$	$\sqrt{26} = 5.0990$
13/2	$7/\sqrt{12} = 2.0207$	$\sqrt{14} = 3.7417$
7	$15/\sqrt{56} = 2.0045$	$\sqrt{30} = 5.4772$
15/2	2.0000	$\sqrt{16} = 4.0000$
8	$17/\sqrt{72} = 2.0035$	$\sqrt{34} = 5.8310$

Table 11: Numerically estimated extreme total splittings ΔE_{min} and ΔE_{max} of $|J\rangle$ states in crystal-fields given in relation to σ .

Quantum number J	ΔE_{min}	ΔE_{max}
1	2.1213	2.4495
3/2	2.0000	2.0000
2	2.0412	3.1623
5/2	2.1213	2.4495
3	2.0207	3.7417
7/2	2.0000	2.8284
4	2.0802	4.0763
9/2	2.1820	3.1623
5	2.2186	3.5904
11/2	2.3075	3.4636
6	2.3350	3.7054
13/2	2.3710	3.7381
7	2.4081	3.8661
15/2	2.3993	3.9880
8	2.3673	4.1043